




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/00	A2	(11) International Publication Number: WO 00/35436 (43) International Publication Date: 22 June 2000 (22.06.00)
(21) International Application Number: PCT/US99/29783 (22) International Filing Date: 15 December 1999 (15.12.99) (30) Priority Data: 60/112,544 16 December 1998 (16.12.98) US 60/164,651 10 November 1999 (10.11.99) US (71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DUDLEY, David, Thomas [US/US]; 3700 Waters Road, Ann Arbor, MI 48103 (US). FLORY, Craig, Mason [US/US]; 3448 Pineview Drive West, Dexter, MI 48130 (US). SALTIEL, Alan, Robert [US/US]; 2002 Valley View Drive, Ann Arbor, MI 48105 (US). (74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.		(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: TREATMENT OF ARTHRITIS WITH MEK INHIBITORS <div style="text-align: center;">  <p>(I)</p> </div> (57) Abstract <p>This invention provides a method for preventing and treating arthritis in a mammal suffering from arthritis and in need of treatment, or suspected of developing arthritis, said method comprising the step of administering an effective antiarthritic amount of a MEK inhibitor, including a compound of Formula (I) and Formula (II).</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TREATMENT OF ARTHRITIS WITH MEK INHIBITORS

FIELD OF THE INVENTION

This invention relates to methods for preventing and treating rheumatoid arthritis or osteoarthritis by administering a compound characterized as an inhibitor of a kinase enzyme known as MEK (MAP kinase or ERK Kinase). MEK phosphorylates and activates MAP kinase (also known as Erk). The method is ideally practiced by administering a phenyl amine MEK inhibitor.

BACKGROUND OF THE INVENTION

Arthritis is a debilitating disease that afflicts millions of people, and for which there currently are no cures. Several forms of arthritis are known. Rheumatoid arthritis is characterized as a chronic systemic inflammatory disease, primarily of the joints, and generally marked by inflammatory changes in the synovial membranes and articular structures and by atrophy and rarefaction of the bones. Osteoarthritis is a noninflammatory degenerative joint disease occurring most often in older persons. Characterized by degeneration of the articular cartilage, hypertrophy of bone at the margins, and changes in the synovial membrane, osteoarthritis is accompanied by pain and stiffness, particularly after prolonged physical activity. Osteoarthritis is also referred to as degenerative arthritis, hypertrophic arthritis, and degenerative joint disease. The current treatments are designed to relieve the pain, and to diminish the symptoms. Most of the known treatments are anti-inflammatory agents such as NSAIDs and cyclooxygenase inhibitors.


We have now discovered that a series of compounds that are said to be selective MEK inhibitors are useful to prevent and treat arthritis. Many of the compounds are described in WO 98/37881 as being useful to treat septic shock.

SUMMARY OF THE INVENTION

This invention provides a method for preventing and treating arthritis, wherein the method comprises the step of administering to a mammal suspected of developing arthritis, or in need of treatment, an effective anti-arthritic amount of a MEK inhibitor, preferably a selective MEK inhibitor. Selective MEK inhibitors are those compounds which inhibit the MEK 1 and MEK 2 enzymes without substantial inhibition of other related enzymes. One aspect of the invention provides a method for treating rheumatoid arthritis, said method comprising the step of administering a MEK inhibitor to a patient. In another aspect, the invention provides a method for treating osteoarthritis, said method comprising administering a MEK inhibitor to a patient. In further embodiments of these aspects, the invention provides a method for preventing and/or treating arthritis comprising the step of administering a therapeutically effective amount of a selective MEK inhibitor described in US 5,525,625, incorporated herein by reference in its entirety. An example of a selective MEK inhibitor is 2-(2-amino-3-methoxyphenyl)-4-oxo-4H-[1]benzopyran.

MEK inhibitors are compounds which inhibit one or more of the family of mammalian enzymes known as MAP kinase kinases, which phosphorylate the MAP kinase subfamily of enzymes (mitogen-associated protein kinase enzymes) referred to as MAP kinases or ERKs (extracellular signal-regulating enzymes such as ERK1 and ERK 2). These enzymes regulate phosphorylation of other enzymes and proteins within the mammalian body. MEK 1 and MEK 2 are dual specificity kinases that are present in all cell types and play a critical role in the regulation of cell proliferation and differentiation in response to mitogens and a wide variety of growth factors and cytokines

In a preferred embodiment, the MEK inhibitor to be administered is a phenyl amine derivative of Formula I



In formula (I), R_1 is hydrogen, hydroxy, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halo, trifluoromethyl, or CN. R_2 is hydrogen. R_3 , R_4 , and R_5 are independently selected from hydrogen, hydroxy, halo, trifluoromethyl, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, nitro, CN, and $-(O \text{ or } NH)_m-(CH_2)_n-R_9$. R_9 is hydrogen, hydroxy, COOH, or $NR_{10}R_{11}$; n is 0-4; m is 0 or 1. Each of R_{10} and R_{11} is independently selected from hydrogen and C_1 - C_8 alkyl, or taken together with the nitrogen to which they are attached can complete a 3-10 member cyclic ring optionally containing 1, 2, or 3 additional heteroatoms selected from O, S, NH, or N- $(C_1$ - C_8 alkyl). Z is $COOR_7$, tetrazolyl, $CONR_6R_7$, $CONHNR_{10}R_{11}$, or CH_2OR_7 . R_6 and R_7 independently are hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, (CO) - C_1 - C_8 alkyl, aryl, heteroaryl, C_3 - C_{10} cycloalkyl, or C_3 - C_{10} (cycloalkyl optionally containing one, two, or three heteroatoms selected from O, S, NH, or N alkyl); or R_6 and R_7 together with the nitrogen to which they are attached complete a 3-10 member cyclic ring optionally containing 1, 2, or 3 additional heteroatoms selected from O, S, NH, or N alkyl. In formula (I), any of the foregoing alkyl, alkenyl, aryl, heteroaryl, heterocyclic, and alkynyl groups can be unsubstituted or substituted by halo, hydroxy, C_1 - C_6 alkoxy, amino, nitro, C_1 - C_4 alkylamino, $di(C_1$ - $C_4)$ alkylamino, C_3 - C_6 cycloalkyl, phenyl, phenoxy, C_3 - C_5 heteroaryl, or C_3 - C_5 heteroaryloxy; or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof.

Preferred embodiments of Formula (I) have a structure wherein: (a) R_1 is hydrogen, methyl, methoxy, fluoro, chloro, or bromo; (b) R_2 is hydrogen; (c) R_3 , R_4 , and R_5 independently are hydrogen, fluoro, chloro, bromo, iodo, methyl, methoxy, or nitro; (d) R_{10} and R_{11} independently are hydrogen or methyl; (e) Z is $COOR_7$, tetrazolyl, $CONR_6R_7$, $CONHNR_{10}R_{11}$, or CH_2OR_7 ; R_6 and R_7

independently are hydrogen, C₁₋₄ alkyl, heteroaryl, or C₃₋₅ cycloalkyl optionally containing one or two heteroatoms selected from O, S, or NH; or R₆ and R₇ together with the nitrogen to which they are attached complete a 5-6 member cyclic ring optionally containing 1 or 2 additional heteroatoms selected from O, NH or N-alkyl; and wherein any of the foregoing alkyl or aryl groups can be unsubstituted or substituted by halo, hydroxy, methoxy, ethoxy, or heteroaryloxy (such as the synthetic intermediate 2,3,4,5,6-pentafluorophenyl); (f) Z is COOR₇; (g) R₇ is H, pentafluorophenyl, or tetrazolyl; (h) R₃, R₄, and R₅ are independently H, fluoro, or chloro; (i) R₄ is fluoro; (j) two of R₃, R₄, and R₅ are fluoro; (k) or combinations of the above. In another preferred embodiment of Formula (I), R₁ is methyl, fluoro, chloro, or bromo.

Examples of preferred embodiments include methods comprising a MEK inhibitor selected from Formula (I) Compound Table below.

FORMULA (I) COMPOUND TABLE
(page 1 of 10)

	[4-Chloro-2-(1H-tetrazol-5-yl)-phenyl-(4-iodo-2-methyl-phenyl)-amine
	(4-iodo-2-methyl-phenyl)-[2-(1H-tetrazol-5-yl)-phenyl]amine
5	[4-nitro-2-(1H-tetrazol-5-yl)-phenyl-(4-iodo-2-methyl-phenyl)-amine
	4-Fluoro-2-(4-iodo-2-methylphenylamino)benzoic acid
	3,4,5-Trifluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
10	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
	Sodium 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoate
	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-benzoic acid
	4-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
15	2-(4-Iodo-2-methyl-phenylamino)-benzoic acid
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
	2,3,5-Trifluoro-4-(4-iodo-2-methyl-phenylamino)-benzoic acid
	2-(4-Iodo-phenylamino)-5-methoxy-benzoic acid
20	5-Methyl-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
	2-(4-Iodo-2-methyl-phenylamino)-4-nitro-benzoic acid
	2-(4-Bromo-2-methyl-phenylamino)-4-fluoro-benzoic acid
	2-(2-Bromo-4-iodo-phenylamino)-5-nitro-benzoic acid
	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-benzoic acid
25	5-Chloro-N-(2-hydroxyethyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-benzamide
	N-Ethyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
30	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(1H-tetrazol-5-yl)- benzamide

FORMULA (I) COMPOUND TABLE
(continued, page 2 of 10)

5	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-benzamide
	[5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoylamino]-acetic acid
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-propyl-benzamide
	5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-
10	benzamide
	N,N-Diethyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	4-Fluoro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo- 2-methyl-phenylamino)-benzamide
	N,N-Diethyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
15	N-Butyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Chloro-N,N-diethyl-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-benzamide
	5-Bromo-3,4-difluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl- phenylamino)-benzamide
20	N-(2,3-Dihydroxy-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)- benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1- yl-ethyl)-benzamide
	3,4-Difluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-
25	benzamide
	N-(2,3-Dihydroxy-propyl)-4-fluoro-2-(4-iodo-2-methyl-phenylamino)- benzamide
	3,4-Difluoro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
30	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin- 1-yl-ethyl)-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyridin-4- yl-ethyl)-benzamide

FORMULA (I) COMPOUND TABLE
(continued, page 3 of 10)

5	4-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Bromo-N-(3-dimethylamino-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide
10	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyridin-4-yl-ethyl)-benzamide
15	N-(3-Dimethylamino-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Benzyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-hydroxy-ethyl)-benzamide
20	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide
25	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-thiophen-2-yl-ethyl)-benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide
30	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-morpholin-4-yl-ethyl)-benzamide

FORMULA (I) COMPOUND TABLE
(continued, page 4 of 10)

5	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide
	2-(4-Bromo-2-methyl-phenylamino)-N-(3-dimethylamino-propyl)-3,4-difluoro-benzamide
10	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyridin-4-yl-ethyl)-benzamide
15	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-pyridin-4-yl-ethyl)-benzamide
	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(3-hydroxy-propyl)-benzamide
	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-pyrrolidin-1-yl-ethyl)-benzamide
20	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-phenethyl-benzamide
	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-thiophen-2-yl-ethyl)-benzamide
	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-pyridin-4-ylmethyl-benzamide
25	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-phenethyl-benzamide
	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-piperidin-1-yl-ethyl)-benzamide
	5-Chloro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide
30	5-Fluoro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide
	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-pyridin-4-yl methyl-benzamide

FORMULA (I) COMPOUND TABLE
(continued, page 5 of 10)

5	5-Bromo-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl- phenylamino)- benzamide
	5-Chloro-N-(2-diethylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)- benzamide
10	(3-Hydroxy-pyrrolidin-1-yl)-[5-nitro-2-(4-iodo-2-methyl-phenylamino)- phenyl]-methanone
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)- benzamide
15	5-Bromo-N-(2-diethylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-chloro-2-(4-iodo-2-methyl- phenylamino)- benzamide
	N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-bromo-2-(4-iodo-2-methyl- phenylamino)- benzamide
20	N-{3-[4-(2-Hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl- phenylamino)- benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl- benzamide
25	5-Bromo-2-(4-iodo-2-ethyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)- benzamide
	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)- benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)- benzamide
30	5-Chloro-N-(3-dimethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-fluoro-2-(4-iodo-2-methyl- phenylamino)- benzamide

FORMULA (I) COMPOUND TABLE
(continued, page 6 of 10)

5	5-Chloro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	5-Chloro-N-(3-diethylamino-2-hydroxy-propyl)-2-(4-iodo-2-methyl- phenylamino)- benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)- benzamide
10	5-Bromo-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)- benzamide
15	N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-2-(4-iodo-2-methyl- phenylamino)-5-nitro- benzamide
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)- benzamide
	5-Chloro-N-(3-diethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
20	5-Chloro-N-(2-diisopropylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)- benzamide
25	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(2-piperidin-1-yl-ethyl)- benzamide
	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperazin-1-yl-ethyl)- benzamide
	N-(2-Diethylamino-ethyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)- benzamide
30	5-Bromo-N-(3-dimethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	N-(3-Hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro- benzamide

FORMULA (I) COMPOUND TABLE
(continued, page 7 of 10)

5	5-Fluoro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-(3-Diethylamino-propyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-(3-Diethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
10	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide
	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(3-piperidin-1-yl-propyl)-benzamide
15	[5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-(2 or 3-hydroxy-pyrrolidin-1-yl)-methanone
	5-Bromo-N-(2-diisopropylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide
20	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide
	[5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-methanone
25	N-(3-Diethylamino-2-hydroxy-propyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)- benzamide
	N-Cyclopropyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
	5-Chloro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide
30	5-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Benzyloxy-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Benzyloxy-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide

FORMULA (I) COMPOUND TABLE
(continued, page 8 of 10)

5	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(4-sulfamoyl-benzyl)- benzamide
	5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	N-(2-Hydroxy-ethyl)-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-(2-Hydroxy-ethyl)-2-(4-iodo-2-ethyl-phenylamino)-5-nitro-benzamide
10	2-(4-Iodo-2-methyl-phenylamino)-N-methyl-5-nitro-N-phenyl-benzamide
	5-Chloro-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl- benzamide
	N-Allyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
15	N-Benzyloxy-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)- benzamide
	N-Allyl-5-chloro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
20	5-Bromo-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl- benzamide
	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)- benzamide
25	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)- benzamide
	N-Allyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(4-sulfamoyl-benzyl)- benzamide
30	N-Allyl-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)- benzamide
	N-Cyclopropyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide


FORMULA (I) COMPOUND TABLE
(continued, page 9 of 10)

5	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl- benzamide
	N-Benzyloxy-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
	N-Cyclohexyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Allyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-benzamide
10	2-(4-Iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-5-nitro-benzamide
	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide
	N-Cyclohexyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Chloro-N-cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-
15	benzamide
	5-Bromo-N-cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-
	benzamide
	N-Cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
20	N-Benzyloxy-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Benzyloxy-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Chloro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-
	benzamide
	5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-
25	benzamide
	2-(4-Iodo-2-methyl-phenylamino)-N-methyl-5-nitro-N-phenyl-benzamide
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-
	benzamide
	N-(2-Hydroxy-ethyl)-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide
30	5-Chloro-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Allyl-5-chloro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-
	benzamide

FORMULA (I) COMPOUND TABLE
(continued, page 10 of 10)

	N-(2-Hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
5	5-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide
	5-Bromo-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Cyclopropyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
10	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)- benzamide
	N-Cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
	N-Allyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Benzyloxy-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide
	N-Allyl-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide
15	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)- benzamide
	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl- benzamide
	N-Allyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide
20	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzyl alcohol
	[5-Chloro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-methanol
	[2-(4-Iodo-2-methyl-phenylamino)-5-nitro-phenyl]-methanol
	[5-Bromo-2-(4-iodo-2-methyl-phenylamino)-phenyl]-methanol
25	N-Allyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide.

In another preferred embodiment, the MEK inhibitor is a compound of
Formula II



In Formula (II), R_{1a} is hydrogen, hydroxy, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halo, trifluoromethyl, or CN. R_{2a} is hydrogen. Each of R_{3a} , R_{4a} , and R_{5a} is independently selected from hydrogen, hydroxy, halo, trifluoromethyl, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, nitro, CN, and $(O \text{ or } NH)_m-(CH_2)_n-R_{9a}$. R_{9a} is hydrogen, hydroxy, CO_2H or $NR_{10a}R_{11a}$; n is 0-4; and m is 0 or 1. Each of R_{10a} and R_{11a} is independently hydrogen or C_1 - C_8 alkyl, or taken together with the nitrogen to which they are attached can complete a 3- to 10-member cyclic ring optionally containing one, two, or three additional heteroatoms selected from O, S, NH, or N- $(C_1$ - C_8 alkyl). R_{6a} is hydrogen, C_1 - C_8 alkyl, $(CO)-(C_1$ - C_8 alkyl), aryl, aralkyl, or C_3 - C_{10} cycloalkyl. R_{7a} is hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, C_3 - C_{10} (cycloalkyl or cycloalkyl optionally containing a heteroatom selected from O, S, or NR_{9a}). In Formula (II), any of the alkyl, alkenyl, aryl, heterocyclic, and alkynyl groups can be unsubstituted or substituted by halo, hydroxy, C_1 - C_6 alkoxy, amino, nitro, C_1 - C_4 alkylamino, di(C_1 - C_4)alkylamino, C_3 - C_6 cycloalkyl, phenyl, phenoxy, C_3 - C_5 heteroaryl, or C_3 - C_5 heteroaryloxy; or R_{6a} and R_{7a} taken together with the N to which they are attached can complete a 5- to 10-membered cyclic ring, optionally containing one, two, or three additional heteroatoms selected from O, S, or $NR_{10a}R_{11a}$. The invention also encompasses pharmaceutically acceptable salts, esters, amides or prodrugs of each of the disclosed compounds.

Preferred embodiments of Formula (II) are those structures wherein:

- (a) R_{1a} is H, methyl, fluoro, or chloro; (b) R_{2a} is H; R_{3a} , R_{4a} , and R_{5a} are each H, Cl, nitro, or F; (c) R_{6a} is H; (d) R_{7a} is methyl, ethyl, 2-propenyl, propyl, butyl, pentyl, hexyl, cyclopropylmethyl, cyclobutylmethyl, cyclopropylmethyl, or cyclopropylethyl;
- 5 (e) the 4' position is I, rather than Br; (f) R_{4a} is F at the 4 position, para to the CO-N- R_{6a} -OR $_{7a}$ group and meta to the bridging nitrogen; (f) R_{3a} or R_{5a} is F; (g) at least one of R_{3a} , R_{4a} , and R_{5a} is F; (h) R_{1a} is methyl or chloro; or (i) or a combination of the above.

10 In a more preferred embodiment the MEK inhibitor is a compound selected from Formula (II) Compound Table below.

FORMULA (II) COMPOUND TABLE
(page 1 of 7)

	4-Fluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide
5	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(methoxy)-benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-ynyloxy)- benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-phenoxyethoxy)- benzamide
10	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-thienylmethoxy)- benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-enyloxy)- benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopropylmethoxy)- benzamide
15	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopentoxo)-benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-furylmethoxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-ethoxy-benzamide
20	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(but-2-enyloxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopropylmethoxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(1-methylprop- 2-ynyloxy)-benzamide
25	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-phenylprop- 2-ynyloxy)-benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl- 5-phenylpent-2-en-4-ynyloxy)-benzamide
30	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-ynyloxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(propoxy)-benzamide

FORMULA (II) COMPOUND TABLE
(continued, page 2 of 7)

5	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclobutyloxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-thienylmethoxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-methyl-prop- 2-enyloxy)-benzamide
10	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-phenoxyethoxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(but-2-enyloxy)- benzamide
15	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(but-3-ynyloxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopentyloxy)- benzamide
	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-(2-fluorophenyl)- prop-2-ynyloxy)-benzamide
20	5-Bromo-3,4-difluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)- benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(n-propoxy)- benzamide
25	5-Bromo-3,4-difluoro-N-(furan-3-ylmethoxy)-2-(4-iodo-2-methyl- phenylamino)-benzamide
	5-Bromo-N-(but-2-enyloxy)-3,4-difluoro-2-(4-iodo-2-methyl- phenylamino)-benzamide
	5-Bromo-N-butoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)- benzamide
30	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-but- 2-enyloxy)-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl- pent-2-en-4-ynyloxy)-benzamide

FORMULA (II) COMPOUND TABLE
(continued, page 3 of 7)

5	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-benzyl)-N-[5-(3-methoxy-phenyl)-3-methyl-pent-2-en-4-ynyloxy]-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-ynyloxy)-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-[3-(3-methoxy-phenyl)-prop-2-ynyloxy]-benzamide
10	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(thiopen-2-ylmethoxy)-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(pyridin-3-ylmethoxy)-benzamide
15	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-(2-fluorophenyl)-prop-2-ynyloxy)-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(ethoxy)-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopropylmethoxy)-benzamide
20	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(isopropoxy)-benzamide
	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-but-3-ynyloxy)-benzamide
	5-Chloro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide
25	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(tetrahydro-pyran-2-yloxy)-benzamide
	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methoxy-benzamide
	4-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-phenylmethoxy-benzamide
	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-phenylmethoxy-benzamide
30	5-Fluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-phenylmethoxy-benzamide
	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(tetrahydropyran-2-yloxy)-benzamide

FORMULA (II) COMPOUND TABLE
(continued, page 4 of 7)

5	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(3-phenylprop-2-ynyloxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(3-furylmethoxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(2-thienylmethoxy)-benzamide
10	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(but-3-ynyloxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(2-methyl-prop-2-ynyloxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(but-2-ynyloxy)-benzamide
15	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(methoxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(ethoxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(cyclobutoxy)-benzamide
20	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(isopropoxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(2-phenoxyethoxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(cyclopropyl-methoxy)-benzamide
25	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(n-propoxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(1-methyl-prop-2-ynyloxy)-benzamide
30	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(3-(3-fluorophenyl)-prop-2-ynyloxy)-benzamide
	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(4,4-dimethylpent-2-ynyloxy)-benzamide

FORMULA (II) COMPOUND TABLE
(continued, page 5 of 7)

5	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(cyclopentoxo)- benzamide
	3,4,5-Trifluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Chloro-3,4-difluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)- benzamide
10	5-Bromo-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy- benzamide
	N-Hydroxy-2-(4-iodo-2-methyl-phenylamino)-4-nitro-benzamide
	3,4,5-Trifluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy-benzamide
	5-Chloro-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy- benzamide
15	5-Bromo-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy- benzamide
	2-(2-Fluoro-4-iodo-phenylamino)-N-hydroxy-4-nitro-benzamide
	2-(2-Chloro-4-iodo-phenylamino)-3,4,5-trifluoro-N-hydroxy-benzamide
	5-Chloro-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy- benzamide
20	5-Bromo-2-(2-bromo-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy- benzamide
	2-(2-Chloro-4-iodo-phenylamino)-N-hydroxy-4-methyl-benzamide
	2-(2-Bromo-4-iodo-phenylamino)-3,4,5-trifluoro-N-hydroxy-benzamide
25	2-(2-Bromo-4-iodo-phenylamino)-5-chloro-3,4-difluoro-N-hydroxy- benzamide
	2-(2-Bromo-4-iodo-phenylamino)-N-hydroxy-4-nitro-benzamide
	4-Fluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy-benzamide
	3,4-Difluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy-benzamide
30	2-(2-Chloro-4-iodo-phenylamino)-4-fluoro-N-hydroxy-benzamide
	2-(2-Chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-benzamide
	2-(2-Bromo-4-iodo-phenylamino)-4-fluoro-N-hydroxy-benzamide
	2-(2-Bromo-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-benzamide

FORMULA (II) COMPOUND TABLE
(continued, page 6 of 7)

5	N-Cyclopropylmethoxy-3,4,5-trifluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Chloro-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide
	5-Bromo-N-cyclopropylmethoxy-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide
10	N-Cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-4-nitro-benzamide
	N-Cyclopropylmethoxy-3,4,5-trifluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide
15	5-Chloro-N-cyclopropylmethoxy-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide
	5-Bromo-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide
	N-Cyclopropylmethoxy-2-(2-fluoro-4-iodo-phenylamino)-4-nitro-benzamide
20	2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4,5-trifluoro-benzamide
	5-Chloro-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide
	5-Bromo-2-(2-bromo-4-iodo-phenylamino)-N-ethoxy-3,4-difluoro-benzamide
25	2-(2-Chloro-4-iodo-phenylamino)-N-ethoxy-4-nitro-benzamide
	2-(2-Bromo-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4,5-trifluoro-benzamide
	2-(2-Bromo-4-iodo-phenylamino)-5-chloro-N-cyclopropylmethoxy-3,4-difluoro-benzamide
30	2-(2-Bromo-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-nitro-benzamide

FORMULA (II) COMPOUND TABLE
(continued, page 7 of 7)

5	N-Cyclopropylmethoxy-4-fluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide
	N-Cyclopropylmethoxy-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide
	2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-fluoro-benzamide
10	2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide
	2-(2-Bromo-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-fluoro-benzamide
15	2-(2-Bromo-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide.

In the most preferred embodiment of this invention, a compound selected from the following is administered to a patient (ie, a mammal) in an amount that is effective to prevent or treat rheumatoid arthritis or osteoarthritis:

2-(2-Chloro-4-iodophenylamino)-N-cyclopropylmethoxy-3,4-difluorobenzamide (PD184352); 2-(2-Methyl-4-iodophenylamino)-N-hydroxy-4-fluorobenzamide (PD170611); 2-(2-Methyl-4-iodophenylamino)-N-hydroxy-3,4-difluoro-5-bromobenzamide (PD171984); 2-(2-Methyl-4-iodophenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-bromobenzamide (PD177168); 2-(2-Methyl-4-iodophenylamino)-N-cyclobutylmethoxy-3,4-difluoro-5-bromobenzamide (PD 180841); 2-(2-Chloro-4-iodophenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-bromobenzamide (PD 184161); 2-(2-Chloro-4-iodophenylamino)-N-hydroxy-3,4-difluoro-5-bromobenzamide (PD184386); 2-(2-Chloro-4-iodophenylamino)-N-cyclobutylmethoxy-3,4-difluorobenzamide (PD 185625); 2-(2-Chloro-4-iodophenylamino)-N-hydroxy-4-fluorobenzamide (PD 185848); 2-(2-Methyl-4-iodophenylamino)-N-hydroxy-3,4-difluorobenzamide (PD 188563); 2-(2-Methyl-4-iodophenylamino)-N-cyclopropylmethoxy-

3,4,5-trifluorobenzamide (PD 198306); and 2-(2-Chloro-4-iodophenylamino)-N-cyclopropylmethoxy-4-fluorobenzamide (PD 203311); and the benzoic acid derivatives thereof. For example, the benzoic acid derivative of PD 198306 is 2-(2-Methyl-4-iodophenylamino)-3,4,5-trifluorobenzoic acid.

5 Additional preferred compounds include 2-(2-chloro-4-iodophenylamino)-5-chloro-N-cyclopropylmethoxy -3,4-difluorobenzamide (PD 297189), 2-(4-iodophenylamino)-N-cyclopropylmethoxy-5-chloro-3,4-difluorobenzamide (PD 297190), 2-(4-iodophenylamino)-5-chloro-3,4-difluorobenzoic acid (PD 296771), 2-(2-chloro-4-iodophenylamino)-5-chloro-10 3,4-difluorobenzoic acid (PD 296770), 5-chloro-3,4-difluoro-2-(4-iodo-2-methylphenylamino)-benzoic acid (PD 296767); and 5-chloro-N-cyclopropylmethoxy -3,4-difluoro-2-(4-iodo-2-methylphenylamino)-benzamide (PD _____).

15 The invention further provides methods of synthesis and synthetic intermediates.

Other features and advantages of the invention are apparent from the detailed description, examples, and claims set forth.

DETAILED DESCRIPTION OF THE INVENTION

20 This invention provides a method of preventing or treating arthritis in a patient which comprises the step of administering to a patient suffering from arthritis and in need of treatment, or to a patient at risk for developing arthritis, an effective anti-arthritic amount of a MEK inhibitor. The invention provides a method of preventing and treating both rheumatoid arthritis and osteoarthritis. The invention is preferably practiced by administering a phenyl amine MEK inhibitor 25 of Formula (I) or Formula (II). Many of these MEK-inhibiting phenyl amine compounds are specific or selective MEK 1 and MEK 2 inhibitors.

30 Selective MEK 1 or MEK 2 inhibitors are those compounds which inhibit the MEK 1 or MEK 2 enzymes without substantially inhibiting other enzymes such as MKK3, ERK, PKC, Cdk2A, phosphorylase kinase, EGF and PDGF receptor kinases, and C-src. In general, a selective MEK 1 or MEK 2 inhibitor has

an IC₅₀ for MEK 1 or MEK 2 that is at least one-fiftieth (1/50) that of its IC₅₀ for one of the above-named other enzymes. Preferably, a selective inhibitor has an IC₅₀ that is at least 1/100, more preferably 1/500, and even more preferably 1/1000, 1/5000 or less than that of its IC₅₀ for one or more of the above-named enzymes.

The mammals to be treated according to this invention are patients, not only humans but also animals such as horses and dogs, who have developed arthritis and are suffering from the pain and disfiguration associated with arthritis, or who are at risk for developing the disease, for example, those who have a family history of arthritis. Those skilled in the medical art are readily able to identify individual patients who are afflicted with arthritis, as well as those who are susceptible to developing the disease.

The term "patient" means all animals including humans. Examples of patients include humans, cows, dogs, cats, goats, sheep, horses, and pigs.

The compounds of the present invention, which can be used to treat septic shock, are MEK inhibitors. A MEK inhibitor is a compound that shows MEK inhibition when tested in the assays titled "Enzyme Assays" in United States Patent Number 5,525,625, column 6, beginning at line 35. The complete disclosure of United States Patent Number 5,525,625 is hereby incorporated by reference. An example of a MEK inhibitor is 2-(2-amino-3-methoxyphenyl)-4-oxo-4H-[1]benzopyran. Specifically, a compound is a MEK inhibitor if a compound shows activity in the assay titled "Cascade Assay for Inhibitors of the MAP Kinase Pathway," column 6, line 36 to column 7, line 4 of the United States Patent Number 5,525,625 and/or shows activity in the assay titled "In Vitro MEK Assay" at column 7, lines 4 to 27 of the above-referenced patent.

A. Terms

Some of the terms used herein are defined below in combination with their usage throughout this disclosure.

As used herein, the term "aryl" means a cyclic, bicyclic, or tricyclic aromatic ring moiety having from five to twelve carbon atoms. Examples of typical aryl groups include phenyl, naphthyl, and fluorenyl. The aryl may be substituted by one, two, or three groups selected from fluoro, chloro, bromo, iodo,

alkyl, hydroxy, alkoxy, nitro, amino, alkylamino, or dialkylamino. Typical substituted aryl groups include 3-fluorophenyl, 3,5-dimethoxyphenyl, 4-nitronaphthyl, 2-methyl-4-chloro-7-aminofluorenyl, and the like.

The term "aryloxy" means an aryl group bonded through an oxygen atom, for example phenoxy, 3-bromophenoxy, naphthyloxy, and 4-methyl-1-fluorenyloxy.

"Heteroaryl" means a cyclic, bicyclic, or tricyclic aromatic ring moiety having from four to eleven carbon atoms and one, two, or three heteroatoms selected from O, S, or N. Examples include furyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, thiazolyl, oxazolyl, xanthenyl, pyronyl, indolyl, pyrimidyl, naphthyridyl, pyridyl, benzinnidazolyl, and triazinyl. The heteroaryl groups can be unsubstituted or substituted by one, two, or three groups selected from fluoro, chloro, bromo, iodo, alkyl, hydroxy, alkoxy, nitro, amino, alkylamino, or dialkylamino. Examples of substituted heteroaryl groups include chloropyranyl, methylthienyl, fluoropyridyl, amino-1,4-benzisoxazinyl, nitroisoquinolyl, and hydroxyindolyl.

The heteroaryl groups can be bonded through oxygen to make heteroaryloxy groups, for example thienyloxy, isothiazolyloxy, benzofuranyloxy, pyridyloxy, and 4-methylisoquinolinyloxy.

The term "alkyl" means straight and branched chain aliphatic groups. Typical alkyl groups include methyl, ethyl, isopropyl, tert.-butyl, 2,3-dimethylhexyl, and 1,1-dimethylpentyl. The alkyl groups can be unsubstituted or substituted by halo, hydroxy, alkoxy, amino, alkylamino, dialkylamino, cycloalkyl, aryl, aryloxy, heteroaryl, or heteroaryloxy, as those terms are defined herein. Typical substituted alkyl groups include chloromethyl, 3-hydroxypropyl, 2-dimethylaminobutyl, and 2-(hydroxymethylamino)ethyl. Examples of aryl and aryloxy substituted alkyl groups include phenylmethyl, 2-phenylethyl, 3-chlorophenylmethyl, 1,1-dimethyl-3-(2-nitrophenoxy)butyl, and 3,4,5-trifluoronaphthylmethyl. Examples of alkyl groups substituted by a heteroaryl or heteroaryloxy group include thienylmethyl, 2-furylethyl, 6-furyloxyoctyl, 4-methylquinolyloxymethyl, and 6-isothiazolylhexyl. Cycloalkyl substituted alkyl groups include cyclopropylmethyl, 2-cyclohexylethyl, piperidyl-2-methyl, 2-(piperidin-1-yl)-ethyl, 3-(morpholin-4-yl)propyl.

“Alkenyl” means a straight or branched carbon chain having one or more double bonds. Examples include but-2-enyl, 2-methyl-prop-2-enyl, 1,1-dimethyl-hex-4-enyl, 3-ethyl-4-methyl-pent-2-enyl, and 3-isopropyl-pent-4-enyl. The alkenyl groups can be substituted with halo, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aryl, aryloxy, heteroaryl, or heteroaryloxy, for example 2-bromoethenyl, 3-hydroxy-2-butenyl, 1-aminoethenyl, 3-phenylprop-2-enyl, 6-thienyl-hex-2-enyl, 2-furyloxy-but-2-enyl, and 4-naphthyloxy-hex-2-enyl.

“Alkynyl” means a straight or branched carbon chain having at least one triple bond. Typical alkynyl groups include prop-2-ynyl, 2-methyl-hex-5-ynyl, 3,4-dimethyl-hex-5-ynyl, and 2-ethyl-but-3-ynyl. The alkynyl groups can be substituted as the alkyl and alkenyl groups, for example, by aryl, aryloxy, heteroaryl, or heteroaryloxy, for example 4-(2-fluorophenyl)-but-3-ynyl, 3-methyl-5-thienylpent-4-ynyl, 3-phenoxy-hex-4-ynyl, and 2-furyloxy-3-methyl-hex-4-ynyl.

The alkenyl and alkynyl groups can have one or more double bonds or triple bonds, respectively, or a combination of double and triple bonds. For example, typical groups having both double and triple bonds include hex-2-en-4-ynyl, 3-methyl-5-phenylpent-2-en-4-ynyl, and 3-thienyloxy-hex-3-en-5-ynyl.

The term “cycloalkyl” means a nonaromatic ring or fused rings. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, bicycloheptyl, adamantyl, and cyclohexyl. The ring can optionally contain one, two, or three heteroatoms selected from O, S, or N. Such groups include tetrahydrofuryl, tetrahydropyrrolyl, octahydrobenzofuranyl, morpholinyl, piperazinyl, pyrrolidinyl, piperidinyl, octahydroindolyl, and octahydrobenzothiofuranyl. The cycloalkyl groups can be substituted with the same substituents as an alkyl and alkenyl groups, for example, halo, hydroxy, aryl, and heteroaryloxy. Examples include 3-hydroxycyclohexyl, 2-aminocyclopropyl, 2-phenylpyrrolidinyl, and 3-thienylmorpholine-1-yl.

B. Administration and Formulation

The MEK inhibitors of the present method can be administered to a patient as part of a pharmaceutically acceptable composition. The compositions can be administered to humans and animals either orally, rectally, parenterally (intravenously, intramuscularly, or subcutaneously), intracisternally,

intravaginally, intraperitoneally, intravesically, locally (powders, ointments, or drops), or as a buccal or nasal spray.

Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents, or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, (c) humectants, as for example, glycerol, (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates, and sodium carbonate, (e) solution retarders, as for example paraffin, (f) absorption accelerators, as for example, quaternary ammonium compounds, (g) wetting agents, as for example, cetyl alcohol and glycerol monostearate, (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid

polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents.

5 Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethyleneglycols, and the like.

10 Solid dosage forms such as tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells, such as enteric coatings and others well-known in the art. They may contain opacifying agents, and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed manner. Examples of embedding compositions which can be used are polymeric substances and waxes. The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

15 Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propyleneglycol, 1,3-butyleneglycol, 20 dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil and sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethyleneglycols, and fatty acid esters of sorbitan or mixtures of these substances, and the like.

25 Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

30 Suspensions, in addition to the active compounds, may contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, or mixtures of these substances, and the like.

Compositions for rectal administrations are preferably suppositories which can be prepared by mixing the compounds of the present invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethyleneglycol, or a

suppository wax, which are solid at ordinary temperatures but liquid at body temperature and therefore, melt in the rectum or vaginal cavity and release the active component.

5 Dosage forms for topical administration of a compound of this invention include ointments, powders, sprays, and inhalants. The active component is admixed under sterile conditions with a physiologically acceptable carrier and any preservatives, buffers, or propellants as may be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

10 The compounds of the present method can be administered to a patient at dosage levels in the range of about 0.1 to about 1000 mg per day. For a normal human adult having a body weight of about 70 kg, a dosage in the range of about 0.01 to about 100 mg per kg of body weight per day is preferable. The specific dosage used, however, can vary. For example, the dosage can depend on a
15 numbers of factors including the requirements of the patient, the severity of the condition being treated, and the pharmacological activity of the compound being used. The determination of optimum dosages for a particular patient is well-known to those skilled in the art.

The compounds of the present method can be administered as
20 pharmaceutically acceptable salts, esters, amides, or prodrugs. The term "pharmaceutically acceptable salts, esters, amides, and prodrugs" as used herein refers to those carboxylate salts, amino acid addition salts, esters, amides, and prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for contact with the tissues of patients without
25 undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "salts" refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared *in situ* during
30 the final isolation and purification of the compounds or by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate,

palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate and laurylsulphonate salts, and the like. These may include cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. (See, for example, S.M. Berge, et al., "Pharmaceutical Salts," *J. Pharm. Sci.*, 1977;66:1-19 which is incorporated herein by reference.)

Examples of pharmaceutically acceptable, non-toxic esters of the compounds of this invention include C₁-C₆ alkyl esters wherein the alkyl group is a straight or branched chain. Acceptable esters also include C₅-C₇ cycloalkyl esters as well as arylalkyl esters such as, but not limited to benzyl. C₁-C₄ alkyl esters are preferred. Esters of the compounds of the present invention may be prepared according to conventional methods.

Examples of pharmaceutically acceptable, non-toxic amides of the compounds of this invention include amides derived from ammonia, primary C₁-C₆ alkyl amines and secondary C₁-C₆ dialkyl amines wherein the alkyl groups are straight or branched chain. In the case of secondary amines the amine may also be in the form of a 5 or 6 membered heterocycle containing one nitrogen atom. Amides derived from ammonia, C₁-C₃ alkyl primary amines and C₁-C₂ dialkyl secondary amines are preferred. Amides of the compounds of the invention may be prepared according to conventional methods.

The term "prodrug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

In addition, the compounds of the present method can exist in unsolvated

as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.


5 Some of the compounds of the present method can exist in different stereoisometric forms by virtue of the presence of chiral centers. It is contemplated that all stereoisometric forms of the compounds as well as mixtures thereof, including racemic mixtures, form part of this invention.

C. Synthesis

The examples presented below are intended to illustrate particular embodiments of the invention and are not intended to limit the scope of the specification, including the claims, in any way. After the priority date of the present disclosure, related syntheses and MEK inhibition data were also published in WO 99/01421 and WO 99/01426, hereby incorporated by reference.

The 2-(4-bromo and 4-iodo phenylamino)-benzoic acid derivatives of Formula (I) can be prepared from commercially available starting materials utilizing synthetic methodologies well-known to those skilled in organic chemistry. A typical synthesis is carried out by reacting a 4-bromo or 4-iodo aniline with a benzoic acid having a leaving group at the 2-position to give a 2-(phenylamino)-benzoic acid. This process is depicted in Scheme 1.

Scheme 1



where L is a leaving group, for example halo such as fluoro.

The reaction of aniline and the benzoic acid derivative generally is accomplished by mixing the benzoic acid with an equimolar quantity or excess of

the aniline in an unreactive organic solvent such as tetrahydrofuran or toluene, in the presence of a base such as lithium diisopropylamide, n-butyl lithium, sodium hydride, triethylamine, and Hunig's base. The reaction generally is carried out at a temperature of about -78°C to about 100°C, and normally is complete within about 2 hours to about 4 days. The product can be isolated by removing the solvent, for example by evaporation under reduced pressure, and further purified, if desired, by standard methods such as chromatography, crystallization, or distillation.


The 2-(phenylamino)-benzoic acid (e.g., Formula I, where R₇ is hydrogen) can be reacted with an organic or inorganic base such as pyridine, triethylamine, calcium carbonate, or sodium hydroxide to produce a pharmaceutically acceptable salt. The free acids can also be reacted with an alcohol of the formula HOR₇ (where R₇ is other than hydrogen, for example methyl) to produce the corresponding ester. Reaction of the benzoic acid with an alcohol can be carried out in the presence of a coupling agent. Typical coupling reagents include 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), 1,3-dicyclohexylcarbodiimide (DCC), bromo-tris(pyrrolidino)- phosphonium hexafluorophosphate (PyBrOP), and (benzotriazolyloxy) tripyrrolidino phosphonium hexafluorophosphate (PyBOP). The phenylamino benzoic acid and alcohol derivative normally are mixed in approximately equimolar quantities in an unreactive organic solvent such as dichloromethane, tetrahydrofuran, chloroform, or xylene, and an equimolar quantity of the coupling reagent is added. A base such as triethylamine or diisopropylethylamine can be added to act as an acid scavenger if desired. The coupling reaction generally is complete after about 10 minutes to 2 hours, and the product is readily isolated by removing the reaction solvent, for instance by evaporation under reduced pressure, and purifying the product by standard methods such as chromatography or crystallizations from solvents such as acetone, diethyl ether, or ethanol.

The benzamides of the invention, Formula (I) where Z is CONR₆R₇, are readily prepared by reacting the foregoing benzoic acids with an amine of the formula HNR₆R₇. The reaction is carried out by reacting approximately equimolar quantities of the benzoic acid and amine in an unreactive organic

solvent in the presence of a coupling reagent. Typical solvents are chloroform, dichloromethane, tetrahydrofuran, benzene, toluene, and xylene. Typical coupling reagents include DCC, EEDQ, PyBrOP, and PyBOP. The reaction is generally complete after about 10 minutes to about 2 hours when carried out at a temperature of about 0°C to about 60°C. The product amide is readily isolated by removing the reaction solvent, for instance by evaporation, and further purification can be accomplished by normal methods such as chromatography, crystallization, or distillation. The hydrazides ($z = \text{CONHNR}_{10}\text{R}_{11}$) are similarly prepared by coupling a benzoic acid with a hydrazine of the formula $\text{H}_2\text{HNR}_{10}\text{R}_{11}$.

The benzyl alcohols of the invention, compounds of Formula (I) where Z is CH_2OR_6 and R_6 is hydrogen, are readily prepared by reduction of the corresponding benzoic acid according to the following Scheme 2.

Scheme 2



Typical reducing agents commonly employed include borane in tetrahydrofuran. The reduction normally is carried out in an unreactive organic solvent such as tetrahydrofuran, and generally is complete within about 2 hours to about 24 hours when conducted at a temperature of about 0°C to about 40°C.

The following detailed examples illustrate specific compounds provided by this invention.

EXAMPLE 1

4-Fluoro-2-(4-iodo-2-methylphenylamino)benzoic acid

To a stirring solution comprised of 3.16 g (0.0133 mol) of 2-amino-5-iodotoluene in 5 mL of tetrahydrofuran at -78°C was added 10 mL (0.020 mol) of a 2.0 M lithium diisopropylamide in tetrahydrofuran/heptane/ethenylbenzene (Aldrich) solution. The resulting green suspension was stirred vigorously for 15 minutes, after which time a solution of 1.00 g (0.00632 mol) of 2,4-difluorobenzoic acid in 10 mL of tetrahydrofuran was added. The reaction temperature was allowed to increase slowly to room temperature, at which temperature it was stirred for 2 days. The reaction mixture was concentrated. Aqueous HCl (10%) was added to the concentrate, and the solution was extracted with dichloromethane. The organic phase was dried (MgSO₄) and then boiled over a steambath to low volume and cooled to room temperature. The off-white fibers were collected by vacuum filtration, rinsed with hexanes, and vacuum-oven dried. (76°C; ca. 10 mm of Hg) to afford 1.10 g (47%) of the desired material; mp 224-229.5°C;

¹H NMR (400 MHz; DMSO): δ 9.72 (s, 1H), 7.97 (dd, 1H, J = 7.0, 8.7 Hz), 7.70 (d, 1H, J = 1.5 Hz), 7.57 (dd, 1H, J = 8.4, 1.9 Hz), 7.17 (d, 1H, J = 8.2 Hz), 6.61-6.53 (m, 2H), 2.18 (s, 3H);

¹³C NMR (100 MHz; DMSO): δ 169.87, 167.60, 165.12, 150.17, 150.05, 139.83, 138.49, 136.07, 135.31, 135.20, 135.07, 125.60, 109.32, 105.09, 104.87, 99.72, 99.46, 89.43, 17.52;

¹⁹F NMR (376 MHz; DMSO): δ -104.00 to -104.07 (m);

IR (KBr) 1670 (C = O stretch) cm⁻¹;

MS (CI) M+1 = 372.

Analysis calculated for C₁₄H₁₁FINO₂:

C, 45.31; H, 2.99; N, 3.77.

Found: C, 45.21; H, 2.77; N, 3.64.

EXAMPLES 2-30

By following the general procedure of Example 1, the following benzoic acids and salts of Formula (I) were prepared.

Example No.	Compound	MP °C
2	3,4,5-Trifluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	206-210
3	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	240.5-244.5
4	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	259.5-262
5	5-Chloro-2-(2-chloro-4-iodo-phenylamino)-benzoic acid	255-260
6	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	234-238
7	Sodium 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoate	310-320 DEC
8	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	239.5-240
9	2-(2-Chloro-4-iodo-phenylamino)-5-nitro-benzoic acid	289-293
10	4-Fluoro-2-(3-fluoro-4-iodo-2-methyl-phenylamino)-benzoic acid	233-235
11	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-benzoic acid	264-267
12	2-(2-Fluoro-4-iodo-phenylamino)-5-nitro-benzoic acid	256-258
13	2-(4-Bromo-2-methyl-phenylamino)-4-fluoro-benzoic acid	218.5-220
14	2-(2-Bromo-4-iodo-phenylamino)-5-nitro-benzoic acid	285-288 DEC
15	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-benzoic acid	230-234
16	3-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	218-221
17	3,4-Difluoro-2-(4-iodo-2-methoxy-phenylamino)-benzoic acid	230-233
18	4-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	245-255 DEC

Example No.	Compound	MP °C
19	2-(4-Iodo-2-methyl-phenylamino)-benzoic acid	218-223
20	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	243-46
21	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	241-245
22	2,3,5-Trifluoro-4-(4-iodo-2-methyl-phenylamino)-benzoic acid	218-222
23	4-Fluoro-2-(3-chloro-4-iodo-2-methyl-phenylamino)-benzoic acid	248-252.5
24	2-(4-Iodo-phenylamino)-5-methoxy-benzoic acid	208-211
25	3-Chloro-2-(2-chloro-4-iodo-phenylamino)-benzoic acid	232-233
26	2-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzoic acid	179-182
27	4-Fluoro-2-(2,3-dimethyl-4-iodo-2-methyl-phenylamino)benzoic acid	258-261
28	5-Methyl-2-(4-iodo-2-methyl-phenylamino)-benzoic acid	209.5-211
29	2-Chloro-6-(4-iodo-2-methyl-phenylamino)-benzoic acid	171-175
30	2-(4-Iodo-2-methyl-phenylamino)-4-nitro-benzoic acid	251-263

EXAMPLE 31

5-Chloro-N-(2-hydroxyethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide

To a stirring solution comprised of 0.1020 g (0.2632 mmol) of 5-chloro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid, 0.1 mL (1.7 mmol) of ethanolamine, and 0.05 mL (0.29 mmol) of diisopropylethylamine in 5 mL of a 1:1 (v/v) tetrahydrofuran-dichloromethane solution was added 0.15 g (0.29 mmol) of solid PyBOP powder directly. The reaction mixture was stirred at room temperature overnight. The solvent was removed in vacuo. The crude residue was partitioned between ether (50 mL) and 10% aqueous hydrochloric acid (50 mL). The organic phase was washed with 10% aqueous sodium hydroxide (50 mL), dried (MgSO₄) and concentrated in vacuo to afford a yellow-brown oil which was crystallized from hexanes-ether to afford 0.0831 g (73%) of a green-yellow powder; mp 120-121°C;

^1H NMR (400 MHz; CDCl_3): δ 9.11 (s, 1H), 7.56 (d, 1H, $J = 1.4$ Hz), 7.46-7.41 (m, 2H), 7.20 (dd, 1H, $J = 8.9, 2.4$ Hz), 7.00 (t, 2H, $J = 9.6$ Hz), 6.55 (broad t, 1H), 3.86 (t, 2H, $J = 5.0$ Hz), 3.61 (dd, 2H, $J = 10.1, 5.5$ Hz), 2.23 (s, 3H), 1.56 (broad s, 1H);

5 IR (KBr) 3297 (O-H stretch), 1627 (C = O stretch) cm^{-1} ;

MS (CI) $M+1 = 431$.

Analysis calculated for $\text{C}_{16}\text{H}_{16}\text{ClIN}_2\text{O}_2$:

C, 44.62; H, 3.74; N, 6.50.

Found: 44.63; H, 3.67; N, 6.30.

10

EXAMPLES 32-48

By following the general procedure of Example 31, the following benzamides were prepared by reacting the corresponding benzoic acid with the corresponding amine.

Example No.	Compound	MP $^{\circ}\text{C}$
32	4-Methoxy-N-(4-methoxy-phenyl)-3-nitro-benzamide	153.5-156
33	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	158
34	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-benzamide	102.5-104.5
35	N-Ethyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	90-91
36	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-benzamide	oil
37	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(1H-tetrazol-5-yl)-benzamide	285-288 DEC
38	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide	180-182
39	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-benzamide	137-138

Example No.	Compound	MP °C
40	[5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoylamino]-acetic acid	170-173
41	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-propyl-benzamide	69-71
42	5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	132-133.4
43	N,N-Diethyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	oil
44	4-Fluoro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide	122-124
45	N,N-Diethyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	91-93
46	N-Butyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	97-99
47	5-Chloro-N,N-diethyl-2-(4-iodo-2-methyl-phenylamino)-benzamide	118-120
48	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-benzamide	142.5-144

EXAMPLE 49

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzyl alcohol

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid (0.50 g, 1.35 mmol) was dissolved in 6 mL (6 mmol) of cold 1.0 M borane-tetrahydrofuran complex in tetrahydrofuran solution. The reaction mixture was stirred under nitrogen atmosphere at room temperature overnight. The reaction was quenched with 80 mL of methanol. Concentration in vacuo produced a clear tan oil which was purified by MPLC. Elution with dichloromethane afforded 0.4285 g (89%) of a white solid; mp 99-100.5°C;

¹H NMR (400 MHz; DMSO): δ 7.57 (d, 1H, J=1.7 Hz), 7.45 (dd, 1H, J=8.4, 1.9 Hz), 7.39 (s, 1H), 7.29 (t, 1H, J=7.5 Hz), 6.89 (d, 1H, J=8.4 Hz), 6.67-6.60 (m, 1H), 5.47 (t, 1H, J=5.5 Hz), 4.49 (d, 2H, 5.1 Hz), 2.14 (s, 3H);

IR (KBr) 3372 (O-H stretch) cm^{-1} ;

MS (CI) $M+1 = 358$.

Analysis calculated for $\text{C}_{14}\text{H}_{13}\text{FINO}$:

C, 47.08; H, 3.67; N, 3.92.

5 Found: C, 47.17; H, 3.75; N, 3.72.

EXAMPLE 50-52

The following benzyl alcohols were prepared by the general procedure of Example 49.

Example No.	Compound	MP °C
50	[5-Chloro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-methanol	82-85
51	[2-(4-Iodo-2-methyl-phenylamino)-5-nitro-phenyl]-methanol	126.5-128.5
52	[5-Bromo-2-(4-iodo-2-methyl-phenylamino)-phenyl]-methanol	60.5-63.5

10 Several invention compounds of Formula (I) were prepared utilizing combinatorial synthetic techniques. The general procedure is as follows:

To a 0.8-mL autosampler vial in a metal block was added 40 μL of a 0.5 M solution of the acid in DMF and 40 μL of the reagent amine (2 M solution in Hunig's base and 1 M in amine in DMF). A 0.5 M solution of PyBrop was freshly prepared and 50 μL were added to the autosampler vial. The reaction was
 15 allowed to stand for 24 hours.

The reaction mixture was transferred to a 2-dram vial and diluted with 2 mL of ethyl acetate. The organic layer was washed with 3 mL of distilled water and the water layer washed again with 2 mL of ethyl acetate. The combined organic layers were allowed to evaporate to dryness in an open fume hood.

20 The residue was taken up in 2 mL of 50% acetonitrile in water and injected on a semi-prep reversed phase column (10 mm \times 25 cm, 5 μM spherical silica, pore size 115 A derivatized with C-18, the sample was eluted at 4.7 mL/min with

a linear ramp to 100% acetonitrile over 8.5 minutes. Elution with 100% acetonitrile continued for 8 minutes). Fractions were collected by monitoring at 214 nM. The residue was dissolved in chloroform and transferred to a preweighed vial, evaporated, and weighed again to determine the yield.

5

EXAMPLES 53-206

The following compounds of Formula I were prepared by combinatorial methodology:

Example No.	Compound	MS M-H
53	5-Bromo-3,4-difluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	510
54	N-(2,3-Dihydroxy-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	462
55	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)-benzamide	577
56	3,4-Difluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	432
57	N-(2,3-Dihydroxy-propyl)-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	444
58	3,4-Difluoro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	446
59	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide	564
60	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyridin-4-yl-ethyl)-benzamide	571
61	4-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	414
62	5-Bromo-N-(3-dimethylamino-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	551

Example No.	Compound	MS M-H
63	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide	580
64	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide	501
65	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide	485
66	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyridin-4-yl-ethyl)-benzamide	493
67	N-(3-Dimethylamino-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	473
68	N-Benzyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	460
69	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-hydroxy-ethyl)-benzamide	384
70	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide	483
71	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide	495
72	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide	513
73	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-thiophen-2-yl-ethyl)-benzamide	480
74	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide	467
75	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-morpholin-4-yl-ethyl)-benzamide	453
76	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide	557
77	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide	479
78	2-(4-Bromo-2-methyl-phenylamino)-N-(3-dimethylamino-propyl)-3,4-difluoro-benzamide	425

Example No.	Compound	MS M-H
79	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide	461
80	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyridin-4-yl-ethyl)-benzamide	475
81	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-pyridin-4-yl-ethyl)-benzamide	445
82	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(3-hydroxy-propyl)-benzamide	400
83	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-pyrrolidin-1-yl-ethyl)-benzamide	437
84	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-phenethyl-benzamide	474
85	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-thiophen-2-yl-ethyl)-benzamide	450
86	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-pyridin-4-ylmethyl-benzamide	431
87	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-phenethyl-benzamide	444
88	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-piperidin-1-yl-ethyl)-benzamide	451
89	5-Chloro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide	557*
90	5-Fluoro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide	541*
91	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-pyridin-4-yl methyl-benzamide	487
92	5-Bromo-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide	601*
93	5-Chloro-N-(2-diethylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	486*

Example No.	Compound	MS M-H
94	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)-benzamide	497*
95	(3-Hydroxy-pyrrolidin-1-yl)-[2-(4-iodo-2-methyl-phenylamino)-5-nitro-phenyl]-methanone	466
96	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide	484*
97	5-Bromo-N-(2-diethylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide	530*
98	N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-chloro-2-(4-iodo-2-methyl- phenylamino)- benzamide	518*
99	N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-bromo-2-(4-iodo-2-methyl- phenylamino)- benzamide	562*
100	[5-Bromo-2-(4-iodo-2-methyl-phenylamino)-phenyl]-(3-hydroxy-pyrrolidin-1-yl)-methanone	499
101	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-benzoic acid phenethyl ester	501
102	N-{3-[4-(2-Hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)- benzamide	568*
103	[5-Chloro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-(3-hydroxy-pyrrolidin-1-yl)-methanone	455
104	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide	460
105	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide	528*
106	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)-benzamide	542*
107	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide	468*
108	5-Chloro-N-(3-dimethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	472*
109	N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-fluoro-2-(4-iodo-2-methyl- phenylamino)- benzamide	502*

Example No.	Compound	MS M-H
110	5-Chloro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	445*
111	5-Chloro-N-(3-diethylamino-2-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide	516*
112	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)-benzamide	482*
113	5-Bromo-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	489*
114	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide	556*
115	N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-2-(4-iodo-2-methyl-phenylamino)-5-nitro- benzamide	529*
116	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide	500*
117	5-Chloro-N-(3-diethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	500*
118	5-Chloro-N-(2-diisopropylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	514*
119	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide	512*
120	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(2-piperidin-1-yl-ethyl)-benzamide	509*
121	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperazin-1-yl-ethyl)-benzamide	544*
122	N-(2-Diethylamino-ethyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	470*
123	5-Bromo-N-(3-dimethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	516*
124	N-(3-Hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	456*
125	5-Fluoro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	429*

Example No.	Compound	MS M-H
126	N-(3-Diethylamino-propyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	484*
127	N-(3-Diethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	511*
128	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide	544*
129	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(3-piperidin-1-yl-propyl)-benzamide	523*
130	[5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-(3-hydroxy-pyrrolidin-1-yl)-methanone	439
131	5-Bromo-N-(2-diisopropylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	558*
132	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide	484*
133	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide	496*
134	[5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-[4-(2-hydroxy-ethyl)-piperazin-1-	482
135	N-(3-Diethylamino-2-hydroxy-propyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)- benzamide	500*
136	[5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoylamino]-acetic acid	443
137	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(2-pyrrolidin-1-yl-ethyl)-benzamide	495*
138	N-(3-Dimethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	483*
139	N-(2-Diisopropylamino-ethyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)- benzamide	498*
140	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-thiobenzoic acid S-phenethyl ester	490
141	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-thiobenzoic acid S-phenethyl ester	506

Example No.	Compound	MS M-H
142	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-thiobenzoic acid S-benzyl ester	536
143	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-thiobenzoic acid S-benzyl ester	503
144	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-thiobenzoic acid S-benzyl ester	476
145	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-thiobenzoic acid S-benzyl ester	492
146	N-Cyclopropyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	409
147	5-Chloro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	429
148	5-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	413
149	N-Benzyloxy-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	475
150	N-Benzyloxy-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide	593*
151	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(4-sulfamoyl-benzyl)-benzamide	567
152	5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	473
153	N-(2-Hydroxy-ethyl)-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide	521

Example No.	Compound	MS M-H
154	N-(2-Hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	440
155	2-(4-Iodo-2-methyl-phenylamino)-N-methyl-5-nitro-N-phenyl-benzamide	486
156	5-Chloro-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-benzamide	425
157	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide	459
158	N-Allyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	409
159	N-Benzoyloxy-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide	583
160	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)-benzamide	538
161	N-Allyl-5-chloro-2-(4-iodo-2-methyl-phenylamino)-benzamide	425
162	N-Cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	436
163	5-Bromo-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-benzamide	469
164	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide	475
165	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)-benzamide	646
166	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)-benzamide	598
167	N-Allyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	436

Example No.	Compound	MS M-H
168	2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(4-sulfamoyl-benzyl)-benzamide	565
169	N-Allyl-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide	469
170	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-benzamide	473
171	N-Cyclopropyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide	517
172	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide	519
173	N-Benzyloxy-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	502
174	N-Cyclohexyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide	559
175	N-Allyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide	517
176	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-benzamide	581
177	2-(4-Iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-5-nitro-benzamide	500
178	5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide	567
179	N-Cyclohexyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	451
180	5-Chloro-N-cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-benzamide	467
181	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-benzamide	533
182	5-Bromo-N-cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-benzamide	511
183	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-benzamide	489
184	N-Cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	478

Example No.	Compound	MS M-H
185	N-Benzyloxy-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide	538
186	N-Benzyloxy-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	477
187	5-Chloro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	431
188	5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	475
189	2-(4-Iodo-2-methyl-phenylamino)-N-methyl-5-nitro-N-phenyl-benzamide	488
190	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide	477
191	N-(2-Hydroxy-ethyl)-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide	523
192	5-Chloro-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-benzamide	425
193	N-Allyl-5-chloro-2-(4-iodo-2-methyl-phenylamino)-benzamide	427
194	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide	461
195	N-(2-Hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	442
196	5-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide	415
197	5-Bromo-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-benzamide	472
198	N-Cyclopropyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	411
199	5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)-benzamide	540
200	N-Cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	438
201	N-Allyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide	411

Example No.	Compound	MS M-H
202	N-Benzyloxy-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide	585
203	N-Allyl-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide	472
204	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)-benzamide	601
205	5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide	522
206	N-Allyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide	438

* M+H

EXAMPLE 207

Preparation of [4-Chloro-2-(1H-tetrazol-5-yl)-(4-iodo-2-methyl-phenyl)-amineStep a: Preparation of 5-chloro-2-fluoro-benzaldehyde

5 To a solution of 1-chloro-4-fluorobenzene (13.06 g, 0.1 mol) in THF (180 mL), at -78°C, LDA (2M solution in THF, 50 mL, 0.1 mol) was added drop wise. After stirring at -78°C for 1.5 hours, DMF (8 mL) was added to the reaction mixture and allowed to warm up to room temperature overnight. The reaction mixture was partitioned between water and Et₂O. The Et₂O layer was dried

10 (MgSO₄) and the solvent removed in vacuum to give 14.95 g (94%) yield of crude aldehyde: ¹H NMR (CDCl₃): δ, 10.3 (s, -C(=O)H).

Step b: Preparation of 5-chloro-2-fluoro-benzaldehyde oxime

A solution of 5-chloro-2-fluoro-benzaldehyde (10 g, 0.0631 mol), hydroxylamine hydrochloride (6.57 g, 0.0946 mol) and pyridine (8.3 mL, 0.1010 mol) in EtOH (100 mL) was heated at 75°C (oil bath temperature) for

15 1 hour and the solvent removed under vacuum to give an oil. The oil was partitioned between water and CH₂Cl₂. The CH₂Cl₂ layer was dried (MgSO₄) and the solvent removed under vacuum to give crude aldoxime as a solid. The solid was purified by medium pressure liquid chromatography on silica. Elution

with CH_2Cl_2 gave 4.87 g (28%) of the aldoxime as white solid: mp 95-97°C;

Analysis calculated for $\text{C}_7\text{H}_5\text{NOFCl}$:

C, 48.44; H, 2.90; N, 8.07.

Found: C, 48.55; H, 2.69, N, 7.90.

5 Step c: Preparation of 5-chloro-2-fluoro-benzonitrile

10 A solution of the 5-chloro-2-fluoro-benzaldehyde oxime (3.15 g, 0.0182 mol) in acetic anhydride (150 mL) was refluxed for 16 hours. The reaction mixture was cooled to room temperature and poured into saturated aqueous NaHCO_3 (200 mL) solution. The mixture was extracted with Et_2O . The Et_2O layer was dried (K_2CO_3) and the solvent removed to give the product as an oily solid. The product was used without further purification in the next step.

Step d: Preparation of 5-(5-chloro-2-fluoro-phenyl)-1H-tetrazole

15 A mixture of 5-chloro-2-fluoro-benzonitrile (2.84 g, 0.01823 mol), butanol (15 mL), sodium azide (1.543 g, 0.0237 mol), acetic acid (1.36 mL, 0.0237 mol) was refluxed for 24 hours. The reaction mixture was cooled to room temperature, additional 1.543 g sodium azide added, and the reaction mixture refluxed for additional 24 hours. After cooling to room temperature, Et_2O (100 mL) and 10% aqueous NaOH (200 mL) were added sequentially. The mixture was vigorously stirred. The aqueous layer was separated, cooled with ice-methanol bath (-15°C) and acidified to pH 1 with conc. HCl . A gray solid precipitated. The solid was dried in vacuum at 50°C to give 1.76 g (49%) of 5-(5-chloro-2-fluoro-phenyl)-1H-tetrazole: mp partial melt at 110°C, complete melting at 124°C);

20 ^1H (400 Mz, CDCl_3): δ 8.19-8.08 (m, 1H), 7.77-7.71 (m, 1H), 7.61-7.52 (m, 1H);

25 ^{13}C (100 Mz, CDCl_3): δ 159.00, 156.49, 140.88, 133.02, 132.93, 130.73, 129.23, 129.21, 129.08, 126.05, 118.96, 118.73, 114.50;

MS (CI) $\text{M}+1 = 199$ (100), $\text{M} = 198$ (6).

Step e: Preparation of [4-Chloro-2-(1H-tetrazol-5-yl)-(4-iodo-2-methyl-phenyl)]-amine

To a solution of 2-methyl-4-iodoaniline (3.52 g, 0.0151 mol) in THF (25 mL) at -78°C, LDA (2 molar solution in THF, 11.33 mL, 0.02267 mol) was added dropwise. After stirring for 0.5 hours, a solution of 1-(tetrazol-5-yl)-2-fluoro-5-chlorobenzene (1.5 g, 0.00756 mol) in THF (15 mL) was added dropwise. The reaction was stirred for 16 hours as it warmed up to room temperature. The reaction mixture was quenched with aqueous conc. NH₄Cl solution and extracted with CH₂Cl₂. The organic layer was dried (MgSO₄) and the solvent removed giving a crude product as an oil. The oil with CH₂Cl₂->CH₂Cl₂:MeOH (9.7:0.3) gave 1.5 g (48%) of the desired product:

mp 205-208°C; ¹H (400 Mz, DMSO): δ 9.13 (s, 1H), 8.00-7.99 (s, 1H), 7.69 (s, 1H), 7.55-7.52 (m, 1H), 7.43-7.40 (m, 1H), 7.12-7.05 (m, 1H), 2.24 (s, 3H); ¹³C (100 Mz, CDCl₃): δ 141.87, 139.28, 138.88, 135.47, 133.71, 131.65, 128.15, 123.69, 121.94, 116.68, 87.79, 17.22; MS (CI) M+2 = 413 (44), M+1 = 412 (85), M = 411 (100).

Analysis calculated for C₁₄H₁₁N₅Cl·0.5H₂O:

C, 39.97; H, 2.87; N, 16.65.

Found: C, 38.87, H, 2.77; N, 16.47.

The following tetrazole substituted phenylamines were prepared by following the general procedure of Example 207.

EXAMPLE 208

(4-iodo-2-methyl-phenyl)-[2-(1H-tetrazol-5-yl)-phenyl]amine, mp 231°C (dec)

EXAMPLE 209


[4-nitro-2-(1H-tetrazol-5-yl)-(4-iodo-2-methyl-phenyl)-amine, mp 205-208°C.

The 4-bromo and 4-iodo phenylamino benzhydroxamic acid derivatives of Formula II can be prepared from commercially available starting materials utilizing synthetic methodologies well-known to those skilled in organic

chemistry. A typical synthesis is carried out by reacting a 4-bromo or 4-iodo aniline with a benzoic acid having a leaving group at the 2-position to give a phenylamino benzoic acid, and then reacting the benzoic acid phenylamino derivative with a hydroxylamine derivative (Scheme 3).

5

Scheme 3



where L is a leaving group, for example halo such as fluoro, chloro, bromo or iodo, or an activated hydroxy group such as a diethylphosphate, trimethylsilyloxy, p-nitrophenoxy, or phenylsulfonyloxy.


5 The reaction of aniline and the benzoic acid derivative generally is accomplished by mixing the benzoic acid with an equimolar quantity or excess of the aniline in an unreactive organic solvent such as tetrahydrofuran, or toluene, in the presence of a base such as lithium diisopropylamide, n-butyl lithium, sodium hydride, and sodium amide. The reaction generally is carried out at a temperature of about -78°C to about 25°C, and normally is complete within about 2 hours to
10 about 4 days. The product can be isolated by removing the solvent, for example by evaporation under reduced pressure, and further purified, if desired, by standard methods such as chromatography, crystallization, or distillation.

The phenylamino benzoic acid next is reacted with a hydroxylamine derivative $\text{HNR}_{6a}\text{OR}_{7a}$ in the presence of a peptide coupling reagent.
15 Hydroxylamine derivatives that can be employed include methoxylamine, N-ethyl-isopropoxy amine, and tetrahydro-oxazine. Typical coupling reagents include 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), 1,3-dicyclohexylcarbodiimide (DCC), bromo-tris(pyrrolidino)-phosphonium hexafluorophosphate (PyBrOP) and (benzotriazolylloxy)tripyrrolidino
20 phosphonium hexafluorophosphate (PyBOP). The phenylamino benzoic acid and hydroxylamino derivative normally are mixed in approximately equimolar quantities in an unreactive organic solvent such as dichloromethane, tetrahydrofuran, chloroform, or xylene, and an equimolar quantity of the coupling reagent is added. A base such as triethylamine or diisopropylethylamine can be
25 added to act as an acid scavenger if desired. The coupling reaction generally is complete after about 10 minutes to 2 hours, and the product is readily isolated by removing the reaction solvent, for instance by evaporation under reduced pressure, and purifying the product by standard methods such as chromatography or crystallizations from solvents such as acetone, diethyl ether, or ethanol.

30 An alternative method for making the invention compounds involves first converting a benzoic acid to a hydroxamic acid derivative, and then reacting the

hydroxamic acid derivative with an aniline. This synthetic sequence is depicted in Scheme 4.


Scheme 4



- 5 where L is a leaving group. The general reaction conditions for both of the steps in Scheme 4 are the same as those described above for Scheme 3.

Yet another method for making invention compounds comprises reacting a phenylamino benzhydroxamic acid with an ester forming group as depicted in Scheme 5.

Scheme 5



where L is a leaving group such as halo, and a base is triethylamine or diisopropylamine.

- 5 The synthesis of compounds of Formula (II) is further illustrated by the following detailed examples.

EXAMPLE 1a

4-Fluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide

(a) Preparation of 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid

- 10 To a stirred solution containing 3.16 g (0.0133 mol) of 2-amino-5-iodotoluene in 5 mL of tetrahydrofuran at -78°C was added 10 mL (0.020 mol) of a 2.0 M lithium diisopropylamide in tetrahydrofuran/heptane/ethylbenzene (Aldrich) solution. The resulting green suspension was stirred vigorously for 15 minutes, after which time a solution of 1.00 g (0.00632 mol) of
- 15 2,4-difluorobenzoic acid in 10 mL of tetrahydrofuran was added. The reaction temperature was allowed to increase slowly to room temperature, at which

temperature the mixture was stirred for 2 days. The reaction mixture was concentrated by evaporation of the solvent under reduced pressure. Aqueous HCl (10%) was added to the concentrate, and the solution was extracted with dichloromethane. The organic phase was dried (MgSO₄) and then concentrated over a steambath to low volume (10 mL) and cooled to room temperature. The off-white fibers which formed were collected by vacuum filtration, rinsed with hexane, and dried in a vacuum-oven (76°C; ca. 10 mm of Hg) to afford 1.10 g (47%) of the desired material; mp 224-229.5°C;

¹H NMR (400 MHz, DMSO): δ 9.72 (s, 1H), 7.97 (dd, 1H, J=7.0, 8.7 Hz), 7.70 (d, 1H, J=1.5 Hz), 7.57 (dd, 1H, J=8.4, 1.9 Hz), 7.17 (d, 1H, J=8.2 Hz), 6.61-6.53 (m, 2H), 2.18 (s, 3H);

¹³C NMR (100 MHz, DMSO): δ 169.87, 166.36 (d, J_{C-F}=249.4 Hz), 150.11 (d, J_{C-F}=11.4 Hz), 139.83, 138.49, 136.07, 135.26 (d, J_{C-F}=11.5 Hz), 135.07, 125.60, 109.32, 104.98 (d, J_{C-F}=21.1 Hz), 99.54 (d, J_{C-F}=26.0 Hz), 89.43, 17.52;

¹⁹F NMR (376 MHz, DMSO): δ -104.00 to -104.07 (m);

IR (KBr) 1670 (C=O stretch)cm⁻¹;

MS (CI) M+1 = 372.

Analysis calculated for C₁₄H₁₁FINO₂:

C, 45.31; H, 2.99; N, 3.77.

Found: C, 45.21; H, 2.77; N, 3.64.

(b) Preparation of 4-Fluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide

To a stirred solution of 4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid (0.6495 g, 0.001750 mol), O-(tetrahydro-2H-pyran-2-yl)-hydroxylamine (0.2590 g, 0.002211 mol), and diisopropylethylamine (0.40 mL, 0.0023 mol) in 31 mL of an equivolume tetrahydrofuran-dichloromethane solution was added 1.18 g (0.00227 mol) of solid PyBOP ([benzotriazolyl]oxy)tripyrrolidino phosphonium hexafluorophosphate, Advanced ChemTech) directly. The reaction mixture was stirred for 30 minutes after which time it was concentrated in vacuo. The brown oil was treated with 10% aqueous hydrochloric acid. The suspension

was extracted with ether. The organic extraction was washed with 10% sodium hydroxide followed by another 10% hydrochloric acid wash, was dried (MgSO₄) and concentrated in vacuo to afford 1.0 g of a light-brown foam. This intermediate was dissolved in 25 mL of ethanolic hydrogen chloride, and the solution was allowed to stand at room temperature for 15 minutes. The reaction mixture was concentrated in vacuo to a brown oil that was purified by flash silica chromatography. Elution with a gradient (100 % dichloromethane to 0.6 % methanol in dichloromethane) afforded 0.2284 g of a light-brown viscous oil. Scratching with pentane-hexanes and drying under high vacuum afforded 0.1541 g (23%) of an off-white foam; mp 61-75°C;

¹H NMR (400 MHz, DMSO): δ 11.34 (s, 1H), 9.68 (s, 1H), 9.18 (s, 1H), 7.65 (d, 1H, J=1.5 Hz), 7.58 (dd, 1H, J=8.7, 6.8 Hz), 7.52 (dd, 1H, J=8.4, 1.9 Hz), 7.15 (d, 1H, J=8.4 Hz), 6.74 (dd, 1H, J=11.8, 2.4 Hz), 6.62 (ddd, 1H, J=8.4, 8.4, 2.7 Hz), 2.18 (s, 3H);

¹³C NMR (100 MHz, DMSO): δ 165.91, 164.36 (d, J_{C-F}=247.1 Hz), 146.78, 139.18, 138.77, 135.43, 132.64, 130.60 (d, J_{C-F}=11.5 Hz), 122.23, 112.52, 104.72 (d, J=22.1 Hz), 100.45 (d, J_{C-F}=25.2 Hz), 86.77, 17.03;

¹⁹F NMR (376 MHz, DMSO): δ -107.20 to -107.27 (m);

IR (KBr) 3307 (broad, O-H stretch), 1636 (C=O stretch) cm⁻¹;

MS (CI) M+1 = 387.

Analysis calculated for C₁₄H₁₂FIN₂O₂:

C, 43.54; H, 3.13; N, 7.25.

Found: C, 43.62; H, 3.24; N, 6.98.

EXAMPLE 2a

25 5-Bromo-3,4-difluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide

(a) Preparation of 5-Bromo-2,3,4-trifluorobenzoic acid

To a stirred solution comprised of 1-bromo-2,3,4-trifluorobenzene (Aldrich, 99%; 5.30 g, 0.0249 mol) in 95 mL of anhydrous tetrahydrofuran cooled to -78°C was slowly added 12.5 mL of 2.0 M lithium diisopropylamide in heptane/tetrahydrofuran/ethylbenzene solution (Aldrich). The mixture was stirred

for 1 hour and transferred by canula into 700 mL of a stirred saturated ethereal carbon dioxide solution cooled to -78°C. The cold bath was removed, and the reaction mixture was stirred for 18 hours at ambient temperature. Dilute (10%) aqueous hydrochloric acid (ca. 500 mL) was poured into the reaction mixture, and the mixture was subsequently concentrated on a rotary evaporator to a crude solid. The solid product was partitioned between diethyl ether (150 mL) and aq. HCl (330 mL, pH 0). The aqueous phase was extracted with a second portion (100 mL) of diethyl ether, and the combined ethereal extracts were washed with 5% aqueous sodium hydroxide (200 mL) and water (100 mL, pH 12). These combined alkaline aqueous extractions were acidified to pH 0 with concentrated aqueous hydrochloric acid. The resulting suspension was extracted with ether (2 × 200 mL). The combined organic extracts were dried (MgSO₄), concentrated in vacuo, and subjected to high vacuum until constant mass was achieved to afford 5.60 g (88% yield) of an off-white powder; mp 139-142.5°C;

¹H NMR (400 MHz, DMSO): δ 13.97 (broad s, 1H, 8.00-7.96 (m, 1H);
¹³C NMR (100 MHz, DMSO): δ 162.96, 129.34, 118.47, 104.54 (d, J_{C-F}=22.9 Hz);
¹⁹F NMR (376 MHz, DMSO): δ -120.20 to -120.31 (m), -131.75 to -131.86 (m), -154.95 to -155.07 (m);
IR (KBr) 1696 (C=O stretch)cm⁻¹;
MS (CI) M+1 = 255.
Analysis calculated for C₇₄H₂₁BrF₃O₂:
C, 32.97; H, 0.79; N, 0.00; Br, 31.34; F, 22.35.
Found: C, 33.18; H, 0.64; N, 0.01; Br, 30.14; F, 22.75.

(b) Preparation of 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid

To a stirred solution comprised of 1.88 g (0.00791 mol) of 2-amino-5-iodotoluene in 10 mL of tetrahydrofuran at -78°C was added 6 mL (0.012 mol) of a 2.0 M lithium diisopropylamide in tetrahydrofuran/heptane/ethylbenzene (Aldrich) solution. The resulting green suspension was stirred vigorously for

10 minutes, after which time a solution of 1.00 g (0.00392 mol) of 5-bromo-2,3,4-trifluorobenzoic acid in 15 mL of tetrahydrofuran was added. The cold bath was subsequently removed, and the reaction mixture stirred for 18 hours. The mixture was concentrated, and the concentrate was treated with 100 mL of dilute
5 (10%) aqueous hydrochloric acid. The resulting suspension was extracted with ether (2 × 150 mL), and the combined organic extractions were dried (MgSO₄) and concentrated in vacuo to give an orange solid. The solid was triturated with boiling dichloromethane, cooled to ambient temperature, and collected by filtration. The solid was rinsed with dichloromethane, and dried in the vacuum-
10 oven (80°C) to afford 1.39 g (76%) of a yellow-green powder; mp 259.5-262°C; ¹H NMR (400 MHz, DMSO): δ 9.03 (s, 1H), 7.99 (dd, 1H, J=7.5, 1.9 Hz), 7.57 (dd, 1H, J=1.5 Hz), 7.42 (dd, 1H, J=8.4, 1.9 Hz), 6.70 (dd, 1H, J=8.4, 6.0 Hz), 2.24 (s, 3H);
¹⁹F NMR (376 MHz, DMSO): δ -123.40 to -123.47 (m); -139.00 to -139.14 (m);
15 IR (KBr) 1667 (C=O stretch)cm⁻¹;
MS (CI) M+1 = 469.

Analysis calculated for C₁₄H₉BrF₂INO₂:

C, 35.93; H, 1.94; N, 2.99; Br, 17.07; F, 8.12; I, 27.11.

Found: C, 36.15; H, 1.91; N, 2.70; Br, 16.40; F, 8.46; I, 26.05.

20 (c) Preparation of 5-Bromo-3,4-difluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide

To a stirred solution comprised of 5-bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid (0.51 g, 0.0011 mol), O-(tetrahydro-2H-pyran-2-yl)-hydroxylamine (0.15 g, 0.0013 mol), and diisopropylethylamine
25 (0.25 mL, 0.0014 mol) in 20 mL of an equivolume tetrahydrofuran-dichloromethane solution was added 0.6794 g (0.001306 mol) of solid PyBOP (Advanced ChemTech) directly. The reaction mixture was stirred at 24°C for 10 minutes, and then was concentrated to dryness in vacuo. The concentrate was suspended in 100 mL of 10% aqueous hydrochloric acid. The suspension was
30 extracted with 125 mL of diethyl ether. The ether layer was separated, washed with 75 mL of 10% aqueous sodium hydroxide, and then with 100 mL of dilute

acid. The ether solution was dried (MgSO_4) and concentrated in vacuo to afford 0.62 g (100%) of an off-white foam. The foam was dissolved in ca. 15 mL of methanolic hydrogen chloride. After 5 minutes, the solution was concentrated in vacuo to an oil, and the oil was purified by flash silica chromatography. Elution with dichloromethane: dichloromethane-methanol (99:1) afforded 0.2233 g (42%) of a yellow powder. The powder was dissolved in diethyl ether and washed with dilute hydrochloric acid. The organic phase was dried (MgSO_4) and concentrated in vacuo to afford 0.200 g of a foam. This product was triturated with pentane to afford 0.1525 g of a powder that was repurified by flash silica chromatography. Elution with dichloromethane afforded 0.0783 g (15%) of an analytically pure title compound, mp 80-90°C;

^1H NMR (400 MHz, DMSO): δ 11.53 (s, 1H), 9.38 (s, 1H), 8.82 (s, 1H), 7.70 (dd, 1H, $J=7.0, 1.9$ Hz), 7.53 (s, 1H), 7.37 (dd, 1H, $J=8.4, 1.9$ Hz), 6.55 (dd, 1H, $J=8.2, 6.5$ Hz), 2.22 (s, 3H);

^{19}F NMR (376 MHz, DMSO): δ -126.24 to -126.29 (m), -137.71 to -137.77 (m);

IR (KBr) 3346 (broad, O-H stretch), 1651 ($\text{C}=\text{O}$ stretch) cm^{-1} ;

MS (CI) $M+1 = 484$.

Analysis calculated for $\text{C}_{14}\text{H}_{10}\text{BrF}_2\text{IN}_2\text{O}_2$:

C, 34.81; H, 2.09; N, 5.80.

Found: C, 34.53; H, 1.73; N, 5.52.

Examples 3a to 12a in the table below were prepared by the general procedure of Examples 1a and 2a.

EXAMPLES 13a-77a

Examples 13a to 77a were prepared utilizing combinatorial synthetic methodology by reacting appropriately substituted phenylamino benzoic acids

(e.g., as shown in Scheme 1) and hydroxylamines (e.g., $(\text{NHR}_{6a})\text{-O-R}_{7a}$). A general method is given below:

To a 0.8-mL autosampler vial in a metal block was added 40 μL of a 0.5 M solution of the acid in DMF and 40 μL of the hydroxylamine (2 M solution

in Hunig's base and 1 M in amine in DMF). A 0.5 M solution of PyBrOP was freshly prepared, and 50 μ L were added to the autosampler vial. The reaction was allowed to stand for 24 hours.

5 The reaction mixture was transferred to a 2-dram vial and diluted with 2 mL of ethyl acetate. The organic layer was washed with 3 mL of distilled water and the water layer washed again with 2 mL of ethyl acetate. The combined organic layers were allowed to evaporate to dryness in an open fume hood.

10 The residue was taken up in 2 mL of 50% acetonitrile in water and injected on a semi-prep reversed phase column (10 mm \times 25 cm, 5 μ M spherical silica, pore Size 115 A derivatized with C-18, the sample was eluted at 4.7 mL/min with a linear ramp to 100% acetonitrile over 8.5 minutes. Elution with 100% acetonitrile continued for 8 minutes.) Fractions were collected by monitoring at 214 nM. The desired fractions were evaporated using a Zymark Turbovap. The product was dissolved in chloroform and transferred to a preweighed vial,
15 evaporated, and weighed again to determine the yield. The structure was confirmed by mass spectroscopy.

EXAMPLES 3a-77a

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
3a	2-(4-bromo-2-methyl-phenylamino)-4-fluoro-N-hydroxy-benzamide	56-75 dec	523
4a	5-Chloro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide	65 dec	
5a	5-Chloro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-N-methyl-benzamide	62-67	
6a	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(terahydropyran-2-yloxy)benzamide	105-108	
7a	5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methoxybenzamide	64-68	
8a	4-Fluoro-N-hydroxy-2-(4-fluoro-2-methyl-phenylamino)-benzamide	119-135	
9a	4-Fluoro-N-hydroxy-2-(2-methyl phenylamino)-benzamide	101-103	
10a	4-Fluoro-2-(4-fluor-2-methyl-phenylamino)-N-(terahydropyran-2-yloxy)benzamide	142-146	
11a	4-Fluoro-N-hydroxy-2-(4-cluoro-2-methyl-phenylamino)-benzamide	133.5-135	

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
12a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-phenylmethoxy-benzamide	107-109.5	
13a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methoxy-benzamide		399
14a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-methoxy-benzamide		417
15a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-methoxy-benzamide		369
16a	2-(4-Bromo-2-methyl-phenylamino)-N-ethoxy-3,4-difluoro-benzamide		342* (M-EtO)
17a	5-Bromo-N-ethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		509
18a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-isopropoxy-benzamide		445
19a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-isopropoxy-benzamide		397
20a	4-Fluoro-N-(furan-3-ylmethoxy)-2-(4-iodo-2-methyl-phenylamino)-benzamide		465

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
21a	3,4-Difluoro-N-(furan-3-ylmethoxy)-2-(4-iodo-2-methyl-phenylamino)-benzamide		483
22a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(furan-3-ylmethoxy)-benzamide		435
23a	5-Bromo-3,4-difluoro-N-(furan-3-ylmethoxy)-2-(4-iodo-2-methyl-phenylamino)-benzamide		561
24a	5-Bromo-N-(but-2-enyloxy)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		536
25a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-ynyloxy)-benzamide		423
26a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-ynyloxy)-benzamide		441
27a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(1-methyl-prop-2-ynyloxy)-benzamide		455
28a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(1-methyl-prop-2-ynyloxy)-benzamide		407
29a	N-(But-3-ynyloxy)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		455

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
30a	2-(4-Bromo-2-methyl-phenylamino)-N-(but-3-ynyloxy)-3,4-difluoro-benzamide		407
31a	5-Bromo-N-(but-3-ynyloxy)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		533
32a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-phenyl-prop-2-ynyloxy)-benzamide		517
33a	3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(3-phenyl-prop-2-ynyloxy)-benzamide		469
34a	3,4-Difluoro-N-[3-(3-fluoro-phenyl)-prop-2-ynyloxy]-2-(4-iodo-2-methyl-phenylamino)-benzamide		535
35a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-[3-(3-fluoro-phenyl)-prop-2-ynyloxy]-benzamide		487
36a	3,4-Difluoro-N-[3-(2-fluoro-phenyl)-prop-2-ynyloxy]-2-(4-iodo-2-methyl-phenylamino)-benzamide		535
37a	5-Bromo-3,4-difluoro-N-[3-(2-fluoro-phenyl)-prop-2-ynyloxy]-2-(4-iodo-2-methyl-phenylamino)-benzamide		613

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
38a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-5-phenyl-pent-2-en-4-ynyloxy)-benzamide		557* *(M+H)
39a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(3-methyl-5-phenyl-pent-2-en-4-ynyloxy)-benzamide		510
40a	N-Ethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		431
41a	2-(4-Bromo-2-methyl-phenylamino)-N-ethoxy-3,4-difluoro-benzamide		383
42a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-propoxy-benzamide		427
43a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-propoxy-benzamide		445
44a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-propoxy-benzamide		397
45a	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-propoxy-benzamide		523
46a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-isopropoxy-benzamide		427

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
47a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-isopropoxy-benzamide		445
48a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-isopropoxy-benzamide		397
49a	5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-isopropoxy-benzamide		523
50a	N-Cyclobutyloxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		457
51a	2-(4-Bromo-2-methyl-phenylamino)-N-cyclobutyloxy-3,4-difluoro-benzamide		409
52a	N-Cyclopentyloxy-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		453
53a	N-Cyclopentyloxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		471
54a	2-(4-Bromo-2-methyl-phenylamino)-N-cyclopentyloxy-3,4-difluoro-benzamide		423
55a	N-Cyclopropylmethoxy-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		439
56a	N-Cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		457

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
57a	2-(4-Bromo-2-methyl-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide		409
58a	5-Bromo-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)		435
59a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-phenoxy-ethoxy)-benzamide		505
60a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-phenoxy-ethoxy)-benzamide		523
61a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-phenoxy-ethoxy)-benzamide		475
62a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(thiophen-2-ylmethoxy)-benzamide		481
63a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(thiophen-2-ylmethoxy)-benzamide		499
64a	2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(thiophen-2-ylmethoxy)-benzamide		451
65a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-methyl-allyloxy)-benzamide		439

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
66a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)- N-(2-methyl-allyloxy)-benzamide		457
67a	2-(4-Bromo-2-methyl-phenylamino)- 3,4-difluoro-N-(2-methyl-allyloxy)-benzamide		410
68a	N-(But-2-enyloxy)-4-fluoro-2-(4-iodo-2-methyl- phenylamino)-benzamide		439
69a	N-(But-2-enyloxy)-3,4-difluoro-2-(4-iodo- 2-methyl-phenylamino)-benzamide		457
70a	2-(4-Bromo-2-methyl-phenylamino)-N-(but- 2-enyloxy)-3,4-difluoro-benzamide		410
71a	3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)- N-(prop-2-ynyloxy)-benzamide		441
72a	N-(But-3-ynyloxy)-3,4-difluoro-2-(4-iodo- 2-methyl-phenylamino)-benzamide		455
73a	2-(4-Bromo-2-methyl-phenylamino)-N- (4,4-dimethyl-pent-2-ynyloxy)-3,4-difluoro- benzamide		449
74a	N-(But-2-enyloxy)-3,4-difluoro-2-(4-iodo- 2-methyl-phenylamino)-benzamide		457

Example No.	Compound	Melting Point (°C)	MS (M-H ⁺)
75a	2-(4-Bromo-2-methyl-phenylamino)-N-(but-2-enyloxy)-3,4-difluoro-benzamide		410
76a	N-(3-tert-butyl-propyn-2-yl)oxy-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide		479
77a	4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-phenylmethoxy-benzamide		577* *CI

D. Pharmacological Activity

Several of the compounds described above have been evaluated in both *in vitro* and *in vivo* assays which are designed to measure anti-arthritic activity, and are recognized by those skilled in the art to be valid predictors of clinical efficacy.

Type II-collagen-induced arthritis (CIA) in mice is recognized as an experimental model of arthritis that has a number of pathologic, immunologic, and genetic features in common with rheumatoid arthritis in humans. The disease is induced by immunization of DBA/1 inbred strain of mice with 100 micrograms of type II collagen (C II), which is the major component of joint cartilage. The collagen was delivered to the mice by intradermal injection of a solution made up in Freund's complete adjuvant. A progressive and inflammatory arthritis develops in the majority of the mice immunized, characterized by paw width increases of up to 100%. A clinical scoring index is used to assess disease progression from erythema and edema (stage 1), joint distortion (stage 2), to joint ankylosis (stage 3). The disease is variable in that it can affect one or all of the paws of the animal, resulting in total possible score of 12 for each mouse. Histopathology of arthritic joints revealed synovitis, pannus formation, and cartilage and bone erosions. All mouse strains that are susceptible to CIA are high antibody responders to type II collagen, and there is a marked cellular response to C II.

The foregoing assay was carried out to evaluate the anti-arthritic activity of several doses of the compound 2-(2-chloro-4-iodophenylamino)-N-cyclopropylmethoxy-3,4-difluorobenzamide. The compound (also referred to as "PD 184352") was suspended in an aqueous mixture of 0.5% hydroxypropylmethyl cellulose (HPMC) and 0.2% Tween 80. The suspension was administered orally twice daily (once in the morning and once in the evening) in equally divided doses. All animals were fed laboratory chow, and given water ad libitum. The assay was continued for 63 days, with disease scores being taken periodically throughout the study, and on Day 63, and averaged at the end of the study. The results of the assay are presented in Pharmacological Table 1 below:

Pharmacological Table 1
Collagen-Induced Arthritis in Mice

Treatment	Number of Mice Per Group	Percent Arthritis Incidence	Average Severity Score	Average No. of Arthritic Paws
Vehicle	10	100	6.3	3.5
PD 184352 200 mg/kg/day	9	22	0.333	0.333
PD 184352 60 mg/kg/day	10	10	0.6	0.4
PD 184352 20 mg/kg/day	10	60	2.9	2

The foregoing data establish that the compound is a potent anti-arthritic agent.

In another standard assay, monoarticular arthritis was induced in rats. Rats were given 6 microgram doses of sonicated Streptococcal cell wall (SCW) in 10 microliters of Dulbecco's phosphate buffered saline (DPBS) by intra-articular injection into the right tibiotalar joint on Day 0. SCW induces paw swelling in the animals. On Day 21, the delayed-type hypersensitivity (DTH) was initiated with 100 micrograms of SCW administered intravenously. Test compounds were suspended in an aqueous mixture of 0.5% HPMC and 0.2% Tween 80, sonicated,

and administered twice daily in equally divided doses (10 mL/kg volume) beginning 1 hour prior to reactivation with SCW. The amount of edema was determined by measuring the baseline volumes of the sensitized hindpaw before reactivation on Day 21, and comparing them with the volumes at subsequent time points. Paw volumes were measured by mercury plethysmography. The antiarthritic activities of several of the phenyl amine compounds described above, when evaluated in the foregoing assay, are presented below in Pharmacological Table 2. In the Table, compound "PD 184352" is 2-(2-chloro-4-iodophenylamino)-N-cyclopropylmethoxy-3,4-difluorobenzamide, and compound "PD 170611" is 2-(2-methyl-4-iodophenylamino)-N-hydroxy-4-fluorobenzamide.

Pharmacological Table 2
SCW-Induced Monoarticular Arthritis

		Percent Inhibition of Paw Swelling at Indicated Time Points			
Treatment	(n)	Day 22	Day 23	Day 24	Day 25
PD 184352 200 mg/kg/day	10	59	57	68	53
PD 184352 60 mg/kg/day	10	30	34	40	31
PD 184352 20 mg/kg/day	10	17	17	28	17
PD 170611 300 mg/kg/day	10	40	4	0	0
PD 170611 100 mg/kg/day	10	80	9	20	32

The foregoing data establish that the phenyl amine compounds of Formulas I and II are potent anti-arthritic agents, and can be used to prevent and

treat various forms of arthritis, including rheumatoid arthritis and osteoarthritis.

Several of the phenyl amine MEK inhibitors have been evaluated in an *in vitro* cell culture assay designed to measure the effect of MEK inhibitors on interleukin-1 (IL-1) induced stromelysin production and phospho-ERK levels in rabbit synovial fibroblasts. The stromelysin is a matrix metalloproteinase enzyme that is a causative factor in arthritis. The phospho-ERK is an enzyme that is phosphorylated by a MEK enzyme, and is thus an indicator of MEK activity in cells.

New England White rabbits were euthanized with B-euthanasia administered IV with a 25 gauge needle in the marginal ear vein. The synovium was immediately removed by the incision of the quadracep tendon and retracting the patella. The synovium, with the infrapellar fat body, was then cut away from the patellar ligament and placed in sterile phosphate buffered saline (PBS) (Gibco BRL, Gaithersberg, MD). The synovium was finely minced with a sterile scalpel and placed in a 50 mL tube containing 6 mL of a solution of 4 mg collagenase type I (Gibco BRL, Gaithersberg, MD)/mL PBS. The mixture was incubated for 3 hours at 37°C. During the incubation, the 50 mL tube was gently swirled 4 to 6 times. The synoviocytes were then washed twice in media (the media composition is described below). Washed cells were seeded into one T-75 plastic cell culture flask and incubated at 37°C in 5% CO₂. After reaching 90-100% confluency, the cells were seeded into appropriate containers for the assay. Synovial fibroblasts were allowed to grow in 96 well plates for three days after confluency before testing. Vehicle (0.1% dimethylsulfoxide in media), or a phenyl amine MEK inhibitor test compound dissolved in vehicle, was added to the synovial fibroblasts 30 minutes before addition of IL-1 α . Interleukin-1 α (100 U/mL) (Genzyme, Cambridge, MA) was suspended in media and added in a volume of 10 μ L/well. The cells were then incubated for 24 hours before the media was removed and stored at -20°C. Prostromelysin-1 levels were measured using an ELISA from Amersham (Cat. No. RPN2615). Percent inhibition was determined by comparing the stromelysin-1 concentration of drug-treated cells to that of vehicle-treated controls. The drug concentration at which 50% inhibition of

stromelysin-1 production was measured (IC₅₀) was determined using linear regression analysis.

The media used in the foregoing assay was prepared as follows, utilizing commercial reagents acquired from Gibco BRL (Gaithersburg, MD) unless otherwise stated. To each 500 mL bottle of alpha-modified Eagles medium (α -MEM, Cat. No. 12561-023) was added 10 mL of 1 Molar N-2-hydroxyethylpiperazine-N-2-ethane sulfonic acid (1 M HEPES, Cat. No. 15630-023), 10 mL of Penicillin/Streptomycin Stock (Cat. No. 15070-030, 5,000 U/mL Pen./5,000 μ g/mL Strep), 500 μ L Gentamicin Stock (50 mg/mL) (Cat. No. 15750-011), 40 mL Fetal Calf Serum from Hyclone Inc. (Cat. No. A1111-L). The results of the foregoing assays are presented in Pharmacological Tables 3 and 4. Pharmacological Table 3 presents the nanomolar dose of test compound required to cause a 50% inhibition of stromelysin expression (IC₅₀).

Pharmacological Table 3

Effect of MEK Inhibitors on IL-1-induced Stromelysin Expression
in Rabbit Synovial Fibroblast Cell Cultures

Compound Tested	IC ₅₀ (nM)
PD 171984	59
PD 177168	20
PD 180841	61
PD 184161	192
PD 184352	28
PD 184386	18
PD 185625	24
PD 185848	9
PD 188563	11
PD 198306	18
PD 199601	24
PD 203311	20

In addition, a Western blot analysis of phospho-ERK levels in cell cultures was performed. Pharmacological Table 4 presents the % inhibition of ERK 1/2 phosphorylation caused by a phenyl amine MEK inhibitor. Cells were lysed with 1mL lysis buffer (containing NaCl (70 mM), B-glycerol phosphate (50 mM), 1M HEPES (10 mM), Triton X-100 (1%)) per T25. The mixture was transferred to microcentrifuge tubes, and spun at 2500 x g for 15 minutes. After removing the supernatant, the protein assay was performed. The samples were run on a 10% Tris-Glycine gel, and transferred to nitrocellulose. The blots were then probed with a phospho-p44/42 MAP kinase antibody followed by the secondary Ab (goat anti rabbit HRP conjugated), coated with the ECL detection reagent, and exposed to film. The amount of phospho-ERK present was determined by relative densitometry.

Pharmacological Table 4

Inhibition of ERK Phosphorylation by PD 184352 in IL-2
Stimulated Rabbit Synovial Fibroblast Cell Cultures

PD 184352 (nM)	% Inhibition of Phospho-ERK
10	22
10	81
1,000	100
10,000	97

The data presented in Pharmacological Tables 3 and 4 establish the phenyl amino MEK inhibitors of Formula I and Formula II are potent inhibitors of cellular enzymes which are causative factors in arthritis.

The method of this invention has also been established in *in vivo* assays utilizing New Zealand White rabbits in which cartilage degradation was induced by interleukin 1-alpha injections into the knee joints.. Adult male rabbits weighing about 3 kg were anesthetized with 5 mg/kg of rompun and 10-15 mg/kg of

ketamine. Test compounds were suspended in a vehicle of 0.5% aqueous hydroxypropyl methyl cellulose and 0.2% Tween 80. The suspensions were administered by oral gavage to the animals. Thirty minutes following dosing with the MEK inhibitors, human recombinant IL-1 α (Genzyme, Cambridge, MA) was injected (25 ng) into one knee joint space through the suprapatella ligament. The contralateral joint received an equal volume of vehicle (phosphate buffered saline/0.2% fetal bovine serum). The animals were euthanized after 24 hours after the IL-1 injection, and the extent of cartilage degradation was determined by measuring the proteoglycan content of the articular cartilage from the femoral condyles with a dimethylene blue dye assay kit. Analysis was done spectrophotometrically, and the percent of inhibition of proteoglycan loss in the treated joint compared to the non-treated joint was determined. The results of this *in vivo* assay for several of the selective MEK inhibitors of Formulas I and II are presented below in Pharmacological Table 5.

Pharmacological Table 5
Inhibition of Proteoglycan Loss in Rabbits

PD No.	Dose (mg/kg)	Dosed	(n)	% Inhibition of Proteoglycan Loss
184352	30	2 \times	6	75
	10	2 \times	6	48
	3	2 \times	6	13
185625	30	1 \times	6	63
185848	30	1 \times	6	43

Additional support for the claimed methods was obtained using the SCW model again and also three other *in vivo* models of inflammation and/or arthritis. The data for each of the additional experiments is shown in Pharmacological Table 6 below. In a carrageenan-induced footpad edema (CFE) model, male outbred Wistar rats (135-150g, Charles River Labs) were dosed orally with 10ml/kg vehicle or test compound one hour prior to administration of a sonicated suspension of carrageenan (1mg/0.1 ml saline). Carrageenan was injected into the

subplantar region of the right hind paw. Paw volume was determined by mercury plethysmography immediately after injection and again five hours after carrageenan injection. Percent inhibition of edema was determined, and the ID₄₀ calculated by linear regression. Differences in swelling compared to control animals were assessed by a 1-way ANOVA, followed by Dunnett's test.

In another model, rat adjuvant-induced polyarthritis (rat AIP) was induced following published procedures. Outbred male Wistar rats (100-115 gms) were obtained from Charles River Labs 2 - 5 days prior to initiation of the study. Rats were injected subcutaneously in the distal third of the tail with 1 mg *Mycobacterium butyricum* suspended in paraffin oil using glass tuberculin syringes and 25 gauge needles on day 0. The *Mycobacterium butyricum* suspension was achieved by sonicating in paraffin oil for 10 minutes with the vessel immersed in an ice bath. After all the rats in the study were immunized, they were randomized into groups. In the therapeutic study, randomization was done on day 14. Dosing started on day 14 and ended on day 27. Vehicle or drug suspended in vehicle was administered orally in 10 ml/kg volume. Hindpaw swelling was assessed by mercury plethysmography, beginning on the 11th day of the study and occurring every third or fourth day subsequently. The change in edema was determined by the difference between hindpaw volume on the day in which it was assessed and the day 14 volume. Percent inhibition was based on a comparison of the treatment group to the vehicle group. The number of animals in a treatment group was 10 while that in the vehicle was 20.

Finally, in a rabbit IL-1 arthritis (IL-1) model, adult male New Zealand White rabbits were anesthetized with rompun (10 mg/kg) and ketamine (50 mg/kg) (im). Twenty-five nanograms IL-1 was injected into one knee joint space through the suprapatella ligament (using sterile techniques). The contralateral joint received an equal volume of vehicle. The knees were first shaved and then swabbed with a surgical disinfectant prior to intraarticular injection. The animals were euthanized after 24 hours, the articular cartilage scraped from the femoral condyles and tibial plateaus and weighed, and the extent of cartilage degradation determined by a standard dimethylene blue assay. Test compound was administered by oral gavage one hour prior to IL-1 administration.

Pharmacological Table 6

Activity of MEK Inhibitors in animal models
of arthritis and inflammation

5


10

Model	184352	198306	203311
Rat carrageenan footpad edema (CFE) (ID ₄₀)	75.8	14.7	18.9 mg/kg
Rat SCW arthritis (SCW) (ID ₅₀)	10.0	11.2	>100 “
Rat adjuvant arthritis (AIP) (ID ₅₀)	6.9	6.6	> 30 “
Rabbit IL-1 arthritis (IL-1) (% Inh proteoglycan loss @ 30mg/kg)	57.9	42.9	29.2

CLAIMS

What is claimed is:

1. A method for treating or preventing arthritis in a mammal, said method comprising the step of administering to a patient suffering from arthritis and in need of treatment, or to a patient suspected of developing arthritis, an effective antiarthritic amount of a MEK inhibitor.
2. The method according to Claim 1 wherein the arthritis is rheumatoid arthritis or osteoarthritis.
3. The method according to Claim 2, wherein the arthritis is osteoarthritis.
4. The method according to Claim 2, wherein the arthritis is rheumatoid arthritis.
5. The method according to Claim 1, wherein said MEK inhibitor is a selective MEK1 or MEK 2 inhibitor.
6. The method according to Claim 1 wherein the MEK inhibitor is a compound of Formula I



wherein:

R_1 is hydrogen, hydroxy, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halo, trifluoromethyl, or CN;

R_2 is hydrogen;

R₃, R₄, and R₅ independently are hydrogen, hydroxy, halo, trifluoromethyl, C₁-C₈ alkyl, C₁-C₈ alkoxy, nitro, CN, or -(O or NH)_m-(CH₂)_n-R₉, where R₉ is hydrogen, hydroxy, COOH, or NR₁₀R₁₁;

5 n is 0-4;

m is 0 or 1;

R₁₀ and R₁₁ independently are hydrogen or C₁-C₈ alkyl, or taken together with the nitrogen to which they are attached can complete a 3-10 member cyclic ring optionally containing 1, 2, or
10 3 additional heteroatoms selected from O, S, NH, or N-C₁-C₈ alkyl;

Z is COOR₇, tetrazolyl, CONR₆R₇, CONHNR₁₀R₁₁, or CH₂OR₇;

R₆ and R₇ independently are hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, (CO)-C₁-C₈ alkyl, aryl, heteroaryl, C₃-C₁₀ cycloalkyl, or
15 C₃-C₁₀ (cycloalkyl optionally containing one, two, or three heteroatoms selected from O, S, NH, or N alkyl); or R₆ and R₇ together with the nitrogen to which they are attached complete a 3-10 member cyclic ring optionally containing 1, 2, or 3 additional heteroatoms selected from O, S, NH, or N alkyl; and wherein any of the foregoing alkyl, alkenyl, aryl,
20 heterocyclic, and alkynyl groups can be unsubstituted or substituted by halo, hydroxy, C₁-C₆ alkoxy, amino, nitro, C₁-C₄ alkylamino, di(C₁-C₄)alkylamino, C₃-C₆ cycloalkyl, phenyl, phenoxy, C₃-C₅ heteroaryl, or C₃-C₅ heteroaryloxy;

or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof.

25 7. The method according to Claim 6 wherein the MEK inhibitor is a compound selected from:

[4-Chloro-2-(1H-tetrazol-5-yl)-phenyl-(4-iodo-2-methyl-phenyl)-amine;

(4-iodo-2-methyl-phenyl)-[2-(1H-tetrazol-5-yl)-phenyl]amine;

- [4-nitro-2-(1H-tetrazol-5-yl)-phenyl-(4-iodo-2-methyl-phenyl)-amine;
- 4-Fluoro-2-(4-iodo-2-methylphenylamino)benzoic acid;
- 3,4,5-Trifluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 5 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- Sodium 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoate;
- 10 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 2-(4-Iodo-2-methyl-phenylamino)-5-nitro-benzoic acid;
- 4-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 2-(4-Iodo-2-methyl-phenylamino)-benzoic acid;
- 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 15 5-Iodo-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 2,3,5-Trifluoro-4-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 2-(4-Iodo-phenylamino)-5-methoxy-benzoic acid;
- 5-Methyl-2-(4-iodo-2-methyl-phenylamino)-benzoic acid;
- 2-(4-Iodo-2-methyl-phenylamino)-4-nitro-benzoic acid;
- 20 2-(4-Bromo-2-methyl-phenylamino)-4-fluoro-benzoic acid;
- 2-(2-Bromo-4-iodo-phenylamino)-5-nitro-benzoic acid;
- 2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-benzoic acid;
- 5-Chloro-N-(2-hydroxyethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 25 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-benzamide;
- N-Ethyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-benzamide;
- 30 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(1H-tetrazol-5-yl)-benzamide;
- 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide;

- 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-
benzamide;
[5-Chloro-2-(4-iodo-2-methyl-phenylamino)-benzoylamino]-acetic
acid;
- 5 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-propyl-benzamide;
5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
N,N-Diethyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
- 10 4-Fluoro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-
2-(4-iodo-2-methyl-phenylamino)-benzamide;
N,N-Diethyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide;
N-Butyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
5-Chloro-N,N-diethyl-2-(4-iodo-2-methyl-phenylamino)-
15 benzamide;
5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N,N-dimethyl-
benzamide;
5-Bromo-3,4-difluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-
phenylamino)-benzamide;
- 20 N-(2,3-Dihydroxy-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-
phenylamino)-benzamide;
5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-
(2-piperidin-1-yl-ethyl)-benzamide;
3,4-Difluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-
25 phenylamino)-benzamide;
N-(2,3-Dihydroxy-propyl)-4-fluoro-2-(4-iodo-2-methyl-
phenylamino)-benzamide;
3,4-Difluoro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-
phenylamino)-benzamide;
- 30 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-
(2-pyrrolidin-1-yl-ethyl)-benzamide;
5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-
(2-pyridin-4-yl-ethyl)-benzamide;

4-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;

5-Bromo-N-(3-dimethylamino-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;

5 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide;

10 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyridin-4-yl-ethyl)-benzamide;

N-(3-Dimethylamino-propyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;

15 N-Benzyl-4-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-hydroxy-ethyl)-benzamide;

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide;

20 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide;

25 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-thiophen-2-yl-ethyl)-benzamide;

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide;

2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-morpholin-4-yl-ethyl)-benzamide;

30 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide;

- 2-(4-Bromo-2-methyl-phenylamino)-N-(3-dimethylamino-propyl)-3,4-difluoro-benzamide;
- 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide;
- 5 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyridin-4-yl-ethyl)-benzamide;
- 2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-pyridin-4-yl-ethyl)-benzamide;
- 2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(3-hydroxy-propyl)-benzamide;
- 10 2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-pyrrolidin-1-yl-ethyl)-benzamide;
- 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-phenethyl-benzamide;
- 15 2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-thiophen-2-yl-ethyl)-benzamide;
- 2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-pyridin-4-ylmethyl-benzamide;
- 2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-phenethyl-benzamide;
- 20 2-(4-Bromo-2-methyl-phenylamino)-3,4-difluoro-N-(2-piperidin-1-yl-ethyl)-benzamide;
- 5-Chloro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 25 5-Fluoro-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-pyridin-4-yl methyl-benzamide;
- 5-Bromo-N-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 30 5-Chloro-N-(2-diethylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;

- 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)-benzamide;
- (3-Hydroxy-pyrrolidin-1-yl)-[2-(4-iodo-2-methyl-phenylamino)-5-nitro-phenyl]-methanone;
- 5 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide;
- 5-Bromo-N-(2-diethylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide;
- N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-chloro-2-(4-iodo-2-methyl- phenylamino)- benzamide;
- 10 N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-bromo-2-(4-iodo-2-methyl- phenylamino)- benzamide;
- N-{3-[4-(2-Hydroxy-ethyl)-piperazin-1-yl]-propyl}-2-(4-iodo-2-methyl-phenylamino)- benzamide;
- 15 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-pyridin-4-ylmethyl-benzamide;
- 5-Bromo-2-(4-iodo-2-ethyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide;
- 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)-benzamide;
- 20 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-pyrrolidin-1-yl-ethyl)-benzamide;
- 5-Chloro-N-(3-dimethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 25 N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-5-fluoro-2-(4-iodo-2-methyl- phenylamino)- benzamide;
- 5-Chloro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5-Chloro-N-(3-diethylamino-2-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)- benzamide;
- 30 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperidin-1-yl-ethyl)-benzamide;

- 5-Bromo-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide;
- 5 N-{2-[Bis-(2-hydroxy-ethyl)-amino]-ethyl}-2-(4-iodo-2-methyl-phenylamino)-5-nitro- benzamide;
- 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide;
- 5-Chloro-N-(3-diethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 10 5-Chloro-N-(2-diisopropylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide;
- 15 2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(2-piperidin-1-yl-ethyl)-benzamide;
- 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-piperazin-1-yl-ethyl)-benzamide;
- N-(2-Diethylamino-ethyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 20 5-Bromo-N-(3-dimethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- N-(3-Hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide;
- 25 5-Fluoro-N-(3-hydroxy-propyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- N-(3-Diethylamino-propyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- N-(3-Diethylamino-propyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide;
- 30 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide;

2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(3-piperidin-1-yl-propyl)-benzamide;

[5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-(3-hydroxy-pyrrolidin-1-yl)-methanone;

5 5-Bromo-N-(2-diisopropylamino-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;

5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-morpholin-4-yl-ethyl)-benzamide;

10 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-piperidin-1-yl-propyl)-benzamide;

[5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-methanone;

N-(3-Diethylamino-2-hydroxy-propyl)-5-fluoro-2-(4-iodo-2-methyl-phenylamino)- benzamide;

15 N-Cyclopropyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;

5-Chloro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;

20 5-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;

N-Benzyloxy-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;

N-Benzyloxy-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide;

25 2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(4-sulfamoyl-benzyl)-benzamide;

5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;

30 N-(2-Hydroxy-ethyl)-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide;

N-(2-Hydroxy-ethyl)-2-(4-iodo-2-ethyl-phenylamino)-5-nitro-benzamide;

- 2-(4-Iodo-2-methyl-phenylamino)-N-methyl-5-nitro-N-phenyl-
benzamide;
- 5-Chloro-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
- 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-
benzamide;
- N-Allyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- N-Benzyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
- 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-
benzyl)-benzamide;
- N-Allyl-5-chloro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- N-Cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-
benzamide;
- 5-Bromo-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
- 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-
benzamide;
- 5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-benzyl)-
benzamide;
- 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-
benzyl)-benzamide;
- N-Allyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide;
- 2-(4-Iodo-2-methyl-phenylamino)-5-nitro-N-(4-sulfamoyl-benzyl)-
benzamide;
- N-Allyl-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-
benzamide;
- N-Cyclopropyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
- 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-
benzamide;

- N-Benzyloxy-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide;
- N-Cyclohexyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5 N-Allyl-5-iodo-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-benzamide;
- 2-(4-Iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-5-nitro-benzamide;
- 10 5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-benzamide;
- N-Cyclohexyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5-Chloro-N-cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 15 5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-benzamide;
- 5-Bromo-N-cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 20 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-benzyl)-benzamide;
- N-Cyclohexyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide;
- N-Benzyloxy-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 25 N-Benzyloxy-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5-Chloro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 30 5-Bromo-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 2-(4-Iodo-2-methyl-phenylamino)-N-methyl-5-nitro-N-phenyl-benzamide;


- 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-
benzamide;
N-(2-Hydroxy-ethyl)-5-iodo-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
5
5-Chloro-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
N-Allyl-5-chloro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-
benzamide;
10
N-(2-Hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-5-nitro-
benzamide;
5-Fluoro-N-(2-hydroxy-ethyl)-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
5-Bromo-N-cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-
15
benzamide;
N-Cyclopropyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-
benzyl)-benzamide;
20
N-Cyclopropyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-
benzamide;
N-Allyl-5-fluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
N-Benzoyloxy-5-iodo-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
25
N-Allyl-5-bromo-2-(4-iodo-2-methyl-phenylamino)-benzamide;
5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-(4-sulfamoyl-
benzyl)-benzamide;
5-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-methyl-N-phenyl-
benzamide;
30
N-Allyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide;
4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-benzyl alcohol;
[5-Chloro-2-(4-iodo-2-methyl-phenylamino)-phenyl]-methanol;
[2-(4-Iodo-2-methyl-phenylamino)-5-nitro-phenyl]-methanol;

[5-Bromo-2-(4-iodo-2-methyl-phenylamino)-phenyl]-methanol;
and
N-Allyl-2-(4-iodo-2-methyl-phenylamino)-5-nitro-benzamide.

5 8. The method of claim 6, wherein the MEK inhibitor is a compound
of Formula (I) wherein (a) R₁ is hydrogen, methyl, methoxy, fluoro,
chloro, or bromo; (b) R₂ is hydrogen; (c) R₃, R₄, and R₅ independently are
hydrogen, fluoro, chloro, bromo, iodo, methyl, methoxy, or nitro; (d) R₁₀
and R₁₁ independently are hydrogen or methyl; (e) Z is COOR₇, tetrazolyl,
CONR₆R₇, CONHNR₁₀R₁₁, or CH₂OR₇; R₆ and R₇ independently are
10 hydrogen, C₁₋₄ alkyl, heteroaryl, or C₃₋₅ cycloalkyl optionally containing
one or two heteroatoms selected from O, S, or NH; or R₆ and R₇ together
with the nitrogen to which they are attached complete a 5-6 member cyclic
ring optionally containing 1 or 2 additional heteroatoms selected from O,
NH or N-alkyl; and wherein any of the foregoing alkyl or aryl groups can
15 be unsubstituted or substituted by halo, hydroxy, methoxy, ethoxy, or
heteroaryloxy; (f) Z is COOR₇; (g) R₇ is H, pentafluorophenyl, or
tetrazolyl; (h) R₃, R₄, and R₅ are independently H, fluoro, or chloro; (i) R₄
is fluoro; (j) two of R₃, R₄, and R₅ are fluoro; or (k) or combinations of the
above.

20 9. The method of claim 8, wherein the MEK inhibitor is a compound
of Formula (I) wherein: Z is COOR₇; R₇ is H, pentafluorophenyl, or
tetrazolyl; R₃ and R₅ are independently H, fluoro, or chloro; and R₄ is
fluoro.

25 10. The method of Claim 1 wherein the MEK inhibitor is a compound
of Formula II



II

wherein:

R_{1a} is hydrogen, hydroxy, C₁-C₈ alkyl, C₁-C₈ alkoxy, halo, trifluoromethyl, or CN;

5 R_{2a} is hydrogen;

R_{3a}, R_{4a}, and R_{5a} independently are hydrogen, hydroxy, halo, trifluoromethyl, C₁-C₈ alkyl, C₁-C₈ alkoxy, nitro, CN, or (O or NH)_m-(CH₂)_n-R_{9a}, where R_{9a} is hydrogen, hydroxy, CO₂H or NR_{10a}R_{11a}.

10 n is 0-4;

m is 0 or 1;

R_{10a} and R_{11a} independently are hydrogen or C₁-C₈ alkyl, or taken together with the nitrogen to which they are attached can complete a 3- to 10-member cyclic ring optionally containing one, two, or three additional heteroatoms selected from O, S, NH, or N-C₁-C₈ alkyl;

15

R_{6a} is hydrogen, C₁-C₈ alkyl, (CO)-C₁-C₈ alkyl, aryl, aralkyl, or C₃-C₁₀ cycloalkyl;

R_{7a} is hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl,

20

C₃-C₁₀ (cycloalkyl or cycloalkyl optionally containing a heteroatom selected from O, S, or NR_{9a});

and wherein any of the foregoing alkyl, alkenyl, aryl, heteroaryl, heterocyclic, and alkynyl groups can be unsubstituted or substituted by halo, hydroxy, C₁-C₆ alkoxy, amino, nitro, C₁-C₄ alkylamino, di(C₁-C₄)alkylamino, C₃-C₆ cycloalkyl, phenyl, phenoxy, C₃-C₅ heteroaryl or

25

heterocyclic radical, or C₃-C₅ heteroaryloxy or heterocyclic radical-oxy; or R_{6a} and R_{7a} taken together with the N to which they are attached can complete a 5- to 10-membered cyclic ring, optionally containing one, two, or three additional heteroatoms selected from O, S, or NR_{10a}R_{11a};

5 or a pharmaceutically acceptable salt, ester, amide or prodrug thereof.

11. The method of Claim 10, comprising a MEK inhibitor having a structure of Formula (II) wherein: (a) R_{1a} is H, methyl, fluoro, or chloro; (b) R_{2a} is H; R_{3a}, R_{4a}, and R_{5a} are each H, Cl, nitro, or F; (c) R_{6a} is H; (d) 10 R_{7a} is methyl, ethyl, 2-propenyl, propyl, butyl, pentyl, hexyl, cyclopropylmethyl, cyclobutyl methyl, cyclopropylmethyl, or cyclopropylethyl; and (e) the 4' position is I, rather than Br.

12. The method of claim 11, comprising a MEK inhibitor having a structure of Formula (II) wherein: R_{4a} is F at the 4 position, para to the CO-N-R_{6a}-OR_{7a} group and meta to the bridging nitrogen; at least one of 15 R_{3a} and R_{5a} is F or Cl; and R_{1a} is methyl or chloro.

13. The method of Claim 10, comprising a MEK inhibitor having a formula selected from:

20 4-Fluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide;

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(methoxy)-benzamide;

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-ynyloxy)-benzamide;

25 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-phenoxyethoxy)-benzamide;

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-thienylmethoxy)-benzamide;

30 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-enyloxy)-benzamide;

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopropylmethoxy)-benzamide;

4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopentoxy)-benzamide;

5 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-furylmethoxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-ethoxy-benzamide;

10 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(but-2-enyloxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopropylmethoxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(1-methylprop-2-ynyloxy)-benzamide;

15 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-phenylprop-2-ynyloxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-5-phenylpent-2-en-4-ynyloxy)-benzamide;

20 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-ynyloxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(propoxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclobutyloxy)-benzamide;

25 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-thienylmethoxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-methyl-prop-2-enyloxy)-benzamide;

30 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(2-phenoxyethoxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(but-2-enyloxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(but-3-ynyloxy)-benzamide;

3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopentyloxy)-benzamide;

5 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-(2-fluorophenyl)-prop-2-ynyloxy)-benzamide;

5-Bromo-3,4-difluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide;

10 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(n-propoxy)-benzamide;

5-Bromo-3,4-difluoro-N-(furan-3-ylmethoxy)-2-(4-iodo-2-methyl-phenylamino)-benzamide;

5-Bromo-N-(but-2-enyloxy)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide

15 5-Bromo-N-butoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;

5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-but-2-enyloxy)-benzamide;

20 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-methyl-pent-2-en-4-ynyloxy)-benzamide;

5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-benzyl)-N-[5-(3-methoxy-phenyl)-3-methyl-pent-2-en-4-ynyloxy]-benzamide;

5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(prop-2-ynyloxy)-benzamide;

25 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-[3-(3-methoxy-phenyl)-prop-2-ynyloxy]-benzamide;

5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(thiopen-2-ylmethoxy)-benzamide;

30 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(pyridin-3-ylmethoxy)-benzamide;

5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(3-(2-fluorophenyl)-prop-2-ynyloxy)-benzamide;

- 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(ethoxy)-benzamide;
- 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(cyclopropylmethoxy)-benzamide;
- 5 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-(isopropoxy)-benzamide;
- 5-Bromo-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-N-but-3-ynyloxy)-benzamide;
- 5-Chloro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 10 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-(tetrahydro-pyran-2-yloxy)-benzamide;
- 5-Chloro-2-(4-iodo-2-methyl-phenylamino)-N-methoxy-benzamide;
- 15 4-Bromo-2-(4-iodo-2-methyl-phenylamino)-N-phenylmethoxy-benzamide;
- 4-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-phenylmethoxy-benzamide;
- 5-Fluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 20 5-Iodo-2-(4-iodo-2-methyl-phenylamino)-N-phenylmethoxy-benzamide;
- 5-Fluoro-2-(4-iodo-2-methyl-phenylamino)-N-(tetrahydropyran-2-yloxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(3-phenylprop-2-ynyloxy)-benzamide;
- 25 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(3-furylmethoxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(2-thienylmethoxy)-benzamide;
- 30 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(but-3-ynyloxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(2-methyl-prop-2-enyloxy)-benzamide;

- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(but-
2-enyloxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(methoxy)-
benzamide;
- 5 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(ethoxy)-
benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-
(cyclobutoxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(isopropoxy)-
10 benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-
(2-phenoxyethoxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(cyclopropyl-
methoxy)-benzamide;
- 15 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(n-propoxy)-
benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-(1-methyl-
prop-2-ynyloxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-
20 (3-(3-fluorophenyl)-prop-2-ynyloxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-
(4,4-dimethylpent-2-ynyloxy)-benzamide;
- 3,4-Difluoro-2-(4-bromo-2-methyl-phenylamino)-N-
(cyclopentoxy)-benzamide;
- 25 3,4,5-Trifluoro-N-hydroxy-2-(4-iodo-2-methyl-phenylamino)-
benzamide;
- 5-Chloro-3,4-difluoro-N-hydroxy-2-(4-iodo-2-methyl-
phenylamino)-benzamide;
- 5-Bromo-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-N-
30 hydroxy-benzamide;
- N-Hydroxy-2-(4-iodo-2-methyl-phenylamino)-4-nitro-benzamide;
- 3,4,5-Trifluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy-
benzamide;

- 5-Chloro-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy-benzamide;
- 5-Bromo-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-benzamide;
- 5 2-(2-Fluoro-4-iodo-phenylamino)-N-hydroxy-4-nitro-benzamide;
- 2-(2-Chloro-4-iodo-phenylamino)-3,4,5-trifluoro-N-hydroxy-benzamide;
- 5-Chloro-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-benzamide;
- 10 5-Bromo-2-(2-bromo-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-benzamide;
- 2-(2-Chloro-4-iodo-phenylamino)-N-hydroxy-4-methyl-benzamide;
- 2-(2-Bromo-4-iodo-phenylamino)-3,4,5-trifluoro-N-hydroxy-benzamide;
- 15 2-(2-Bromo-4-iodo-phenylamino)-5-chloro-3,4-difluoro-N-hydroxy-benzamide;
- 2-(2-Bromo-4-iodo-phenylamino)-N-hydroxy-4-nitro-benzamide;
- 4-Fluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy-benzamide;
- 20 3,4-Difluoro-2-(2-fluoro-4-iodo-phenylamino)-N-hydroxy-benzamide;
- 2-(2-Chloro-4-iodo-phenylamino)-4-fluoro-N-hydroxy-benzamide;
- 2-(2-Chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-benzamide;
- 25 2-(2-Bromo-4-iodo-phenylamino)-4-fluoro-N-hydroxy-benzamide;
- 2-(2-Bromo-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-benzamide;
- N-Cyclopropylmethoxy-3,4,5-trifluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 30 5-Chloro-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide;
- 5-Bromo-N-cyclopropylmethoxy-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide;

N-Cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-4-nitro-benzamide;

N-Cyclopropylmethoxy-3,4,5-trifluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide;

5 5-Chloro-N-cyclopropylmethoxy-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide;

5-Bromo-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide;

10 N-Cyclopropylmethoxy-2-(2-fluoro-4-iodo-phenylamino)-4-nitro-benzamide;

2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4,5-trifluoro-benzamide;

5-Chloro-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide;

15 5-Bromo-2-(2-bromo-4-iodo-phenylamino)-N-ethoxy-3,4-difluoro-benzamide;

2-(2-Chloro-4-iodo-phenylamino)-N-ethoxy-4-nitro-benzamide;

2-(2-Bromo-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4,5-trifluoro-benzamide;

20 2-(2-Bromo-4-iodo-phenylamino)-5-chloro-N-cyclopropylmethoxy-3,4-difluoro-benzamide

2-(2-Bromo-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-nitro-benzamide;

25 N-Cyclopropylmethoxy-4-fluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide;

N-Cyclopropylmethoxy-3,4-difluoro-2-(2-fluoro-4-iodo-phenylamino)-benzamide;

2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-fluoro-benzamide;

30 2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide;

2-(2-Bromo-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-fluoro-benzamide; and

2-(2-Bromo-4-iodo-phenylamino)-N-cyclopropylmethoxy-
3,4-difluoro-benzamide.

14. The method of claim 1, comprising a MEK inhibitor having a
structure selected from:

5 2-(2-chloro-4-iodophenylamino)-5-chloro-N-cyclopropylmethoxy -3,4-
difluorobenzamide (PD 297189); 2-(4-iodophenylamino)-N-
cyclopropylmethoxy-5-chloro-3,4-difluorobenzamide (PD 297190); 2-(4-
iodophenylamino)-5-chloro-3,4-difluorobenzoic acid (PD 296771); 2-(2-
chloro-4-iodophenylamino)-5-chloro-3,4-difluorobenzoic acid
10 (PD 296770); 5-chloro-3,4-difluoro-2-(4-iodo-2-methylphenylamino)-
benzoic acid (PD 296767); and 5-chloro-N-cyclopropylmethoxy -3,4-
difluoro-2-(4-iodo-2-methylphenylamino)-benzamide (PD _____).

15 15. A method of treating or preventing arthritis in a patient in need of
treatment, or suspected of developing arthritis, said method comprising the
step of administering an effective antiarthritic amount of a compound
selected from:

2-(2-Chloro-4-iodophenylamino)-N-cyclopropylmethoxy-
3,4-difluorobenzamide (PD184352);
2-(2-Methyl-4-iodophenylamino)-N-hydroxy-4-fluorobenzamide
20 (PD170611);
2-(2-Methyl-4-iodophenylamino)-N-hydroxy-3,4-difluoro-
5-bromobenzamide (PD171984);
2-(2-Methyl-4-iodophenylamino)-N-cyclopropylmethoxy-
3,4-difluoro-5-bromobenzamide (PD177168);
25 2-(2-Methyl-4-iodophenylamino)-N-cyclobutylmethoxy-
3,4-difluoro-5-bromobenzamide (PD 180841);
2-(2-Chloro-4-iodophenylamino)-N-cyclopropylmethoxy-
3,4-difluoro-5-bromobenzamide (PD 184161);
2-(2-Chloro-4-iodophenylamino)-N-hydroxy-3,4-difluoro-
30 5-bromobenzamide (PD184386);

2-(2-Chloro-4-iodophenylamino)-N-cyclobutylmethoxy-
3,4-difluorobenzamide (PD 185625);
2-(2-Chloro-4-iodophenylamino)-N-hydroxy-4-fluorobenzamide
(PD 185848);

5 2-(2-Methyl-4-iodophenylamino)-N-hydroxy-
3,4-difluorobenzamide (PD 188563);
2-(2-Methyl-4-iodophenylamino)-N-cyclopropylmethoxy-
3,4,5-trifluorobenzamide (PD 198306); and
2-(2-Chloro-4-iodophenylamino)-N-cyclopropylmethoxy-
10 4-fluorobenzamide (PD 203311).

16. The method of Claim 15 wherein said compound is selected from
2-(2-chloro-4-iodophenylamino)-N-cyclopropylmethoxy-
3,4-difluorobenzamide (PD 184352), 2-(2-Methyl-4-iodophenylamino)-N-
15 cyclopropylmethoxy-3,4,5-trifluorobenzamide (PD 198306); and
2-(2-Chloro-4-iodophenylamino)-N-cyclopropylmethoxy-
4-fluorobenzamide (PD 203311).

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
22 June 2000 (22.06.2000)

PCT

(10) International Publication Number
WO 00/35436 A3(51) International Patent Classification⁷: **A61K 31/195**,
31/166, 31/136, 31/41, 31/445, 31/495, 31/535, 31/38,
31/34Alan, Robert [US/US]; 2002 Valley View Drive, Ann
Arbor, MI 48105 (US).

(21) International Application Number: PCT/US99/29783

(74) Agents: **RYAN, M., Andrea**; Warner-Lambert Company,
201 Tabor Road, Morris Plains, NJ 07950 et al. (US).(22) International Filing Date:
15 December 1999 (15.12.1999)(81) Designated States (*national*): AE, AL, AU, BA, BB, BG,
BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID,
IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK,
MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA,
US, UZ, VN, YU, ZA.

(25) Filing Language: English


(26) Publication Language: English

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).(30) Priority Data:
60/112,544 16 December 1998 (16.12.1998) US
60/164,651 10 November 1999 (10.11.1999) US(71) Applicant (*for all designated States except US*):
WARNER-LAMBERT COMPANY [US/US]; 201
Tabor Road, Morris Plains, NJ 07950 (US).Published:
— with international search report

(72) Inventors; and

(88) Date of publication of the international search report:
18 October 2001(75) Inventors/Applicants (*for US only*): **DUDLEY, David**,
Thomas [US/US]; 3700 Waters Road, Ann Arbor, MI
48103 (US). **FLORY, Craig, Mason** [US/US]; 3448
Pineview Drive West, Dexter, MI 48130 (US). **SALTIEL,***For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: TREATMENT OF ARTHRITIS WITH MEK INHIBITORS



(I)

(57) Abstract: This invention provides a method for preventing and treating arthritis in a mammal suffering from arthritis and in need of treatment, or suspected of developing arthritis, said method comprising the step of administering an effective antiarthritic amount of a MEK inhibitor, including a compound of Formula (I) and Formula (II).

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/29783

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/195 A61K31/166 A61K31/136 A61K31/41 A61K31/445
A61K31/495 A61K31/535 A61K31/38 A61K31/34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. REFERENCES CONSIDERED TO BE RELEVANT

	ation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P	<p>WO 99 61440 A (SMITHKLINE BEECHAM CO) 2 December 1999 (1999-12-02) claims 1-8</p> <p style="text-align: center;">---</p> <p>WO 99 57101 A (F. HOFFMANN-LA ROCHE AG) 11 November 1999 (1999-11-11) page 52, line 14 - line 23 claims 1,37-43</p> <p style="text-align: center;">---</p> <p>WO 99 01449 A (NOVARTIS AG) 14 January 1999 (1999-01-14) claims 1-6,8-11 page 20, line 16 - line 21 page 23, line 14 -page 24, line 14</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	<p>1-5</p> <p>1-5</p> <p>1-5</p>

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

5 April 2000

Date of mailing of the international search report

19/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Siatou, E

INTERNATIONAL SEARCH REPORT

Int'l. Jonal Application No

PCT/US 99/29783

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	K. MIYAZAWA ET AL: "Regulation of Interleukin-1-beta-induced Interleukin-6 Gene Expression in Human Fibroblast-like Synoviocytes by p38 Mitogen-activated Protein Kinase" THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 38, 18 September 1998 (1998-09-18), pages 24832-24838, XP002134930 abstract	1-5
Y	page 24837, right-hand column, last paragraph	6-16
X	--- J. R. JACKSON ET AL: "Pharmacological Effects of SB 220025, a Selective Inhibitor of P38 Mitogen-Activated Protein Kinase, in Angiogenesis and Chronic Inflammatory Disease Models" THE JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS, vol. 284, no. 2, 1998, pages 687-692, XP002134931 abstract	1-5
Y	page 690, right-hand column, line 7 - line 24	6-16
X	--- A. M. BADGER ET AL: "SB 203580 Inhibits p38 Mitogen-Activated Protein Kinase, Nitric Oxide Production, and iNducible Nitric Oxide Synthase in Bovine Cartilage-Derived Chondrocytes" THE JOURNAL OF IMMUNOLOGY, vol. 161, 1998, pages 467-473, XP002134932 abstract	1-5
Y	page 472, right-hand column, last paragraph	6-16
Y	--- WO 98 37881 A (WARNER LAMBERT COMPANY) 3 September 1998 (1998-09-03) cited in the application claims 1-9	6-16
X	--- HENRY J R ET AL: "Potent inhibitors of the map kinase p38" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, GB, OXFORD, vol. 8, no. 23, 1 December 1998 (1998-12-01), pages 3335-3340, XP004143754 ISSN: 0960-894X abstract page 3336; table 1 page 3337; table 2 --- -/--	1-5

INTERNATIONAL SEARCH REPORT

Int'l. Patent Application No
PCT/US 99/29783

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 28292 A (SMITHKLINE BEECHAM CO) 2 July 1998 (1998-07-02) claims 1-12,16-19 ----	1-5
X	WO 96 00082 A (CORTECS LIMITED) 4 January 1996 (1996-01-04) claims 1-6,12,14,15,20 page 7, line 17 -page 8, line 25 page 11, line 21 ----	1-5
A	US 5 525 625 A (A. J. BRIDGES ET AL) 11 June 1996 (1996-06-11) cited in the application the whole document -----	1-16

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/29783

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-16
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-16
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/29783

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9961440	A	02-12-1999	NONE		
WO 9957101	A	11-11-1999	NONE		
WO 9901449	A	14-01-1999	AU	8801598 A	25-01-1999
			NO	996429 A	23-12-1999
			ZA	9805656 A	30-12-1998
WO 9837881	A	03-09-1998	AU	5610398 A	18-09-1998
			ZA	9801578 A	02-09-1998
WO 9828292	A	02-07-1998	NONE		
WO 9600082	A	04-01-1996	AU	2749395 A	19-01-1996
			CA	2193654 A	04-01-1996
			CN	1151119 A	04-06-1997
			EP	0766565 A	09-04-1997
			FI	965204 A	21-02-1997
			JP	10502073 T	24-02-1998
			NO	965564 A	24-02-1997
US 5525625	A	11-06-1996	AT	181913 T	15-07-1999
			AU	690400 B	23-04-1998
			AU	4245696 A	14-08-1996
			CA	2208075 A	01-08-1996
			DE	69510696 D	12-08-1999
			DE	69510696 T	23-12-1999
			EP	0805807 A	12-11-1997
			GR	3031295 T	31-12-1999
			JP	10512878 T	08-12-1998
			NZ	297320 A	28-05-1999
			WO	9622985 A	01-08-1996
			ZA	9600528 A	15-08-1996